

## **Product datasheet for SC200577**

## 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com

OriGene Technologies, Inc.

https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Dectin 1 (CLEC7A) (NM\_197954) Human 3' UTR Clone

**Product data:** 

**Product Type:** 3' UTR Clones

Product Name: Dectin 1 (CLEC7A) (NM\_197954) Human 3' UTR Clone

Symbol: Dectin 1

**Synonyms:** BGR; CANDF4; CD369; CLECSF12; DECTIN1; SCARE2

Mammalian Cell

Selection:

Neomycin

**Vector:** pMirTarget (PS100062)

**ACCN:** NM\_197954

**Insert Size:** 118 bp

Insert Sequence: >SC200577 3'UTR clone of NM\_197954

The sequence shown below is from the reference sequence of NM\_197954. The complete

sequence of this clone may contain minor differences, such as SNPs.

Blue=Stop Codon Red=Cloning site

GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAGGCCAAGAAGGGCGGAAAGATCGCCGTG

TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC

GGTTTCAAAGCTGTGGAATTCAAAGGATAAATTAATGAAGAAAACAAGCGGAGCTGAAGAAGAAAGTAC

AATATGGTGCTGTCTTCCTAATGAAATAAATTCACTAAATGGACATTAA

**ACGCGT**AAGCGGCCGCGCATCTAGATTCGAAGAAAATGACCGACCAAGCGACGCCCAACCTGCCATCA

CGAGATTTCGATTCCACCGCCGCCTTCTATGAAAGG

**Restriction Sites:** Sgfl-Mlul

OTI Disclaimer: Our molecular clone sequence data has been matched to the sequence identifier above as a

point of reference. Note that the complete sequence of this clone is largely the same as the

reference sequence but may contain minor differences, e.g., single nucleotide

polymorphisms (SNPs).

**Components:** The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The

package also includes 100 pmols of both the corresponding 5' and 3' vector primers in

separate vials.

**RefSeq:** <u>NM 197954.3</u>





## Dectin 1 (CLEC7A) (NM\_197954) Human 3' UTR Clone - SC200577

Summary: This gene encodes a member of the C-type lectin/C-type lectin-like domain (CTL/CTLD)

superfamily. The encoded glycoprotein is a small type II membrane receptor with an

extracellular C-type lectin-like domain fold and a cytoplasmic domain with an

immunoreceptor tyrosine-based activation motif. It functions as a pattern-recognition receptor that recognizes a variety of beta-1,3-linked and beta-1,6-linked glucans from fungi and plants, and in this way plays a role in innate immune response. Alternate transcriptional splice variants, encoding different isoforms, have been characterized. This gene is closely linked to other CTL/CTLD superfamily members on chromosome 12p13 in the natural killer

gene complex region. [provided by RefSeq, Jul 2008]

**Locus ID:** 64581

MW: 4.7