

## **Product datasheet for SC200626**

## PKN1 (NM 002741) Human 3' UTR Clone

## **Product data:**

**Product Type:** 3' UTR Clones

Product Name: PKN1 (NM 002741) Human 3' UTR Clone

Symbol: PKN'

Synonyms: DBK; PAK-1; PAK1; PKN; PKN-ALPHA; PRK1; PRKCL1

Mammalian Cell

Selection:

Neomycin

**Vector:** pMirTarget (PS100062)

**ACCN:** NM\_002741

**Insert Size:** 105 bp

Insert Sequence: >SC200626 3'UTR clone of NM\_002741

The sequence shown below is from the reference sequence of NM\_002741. The complete

sequence of this clone may contain minor differences, such as SNPs.

Blue=Stop Codon Red=Cloning site

GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAGGCCAAGAAGGGCGGAAAGATCGCCGTG

TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC

GACTTCGACTTCGTGGCCGGGGGCTGCTAGCCCCCTCCCCTGCCCCTGCCCCTGCCCCGAGAGC

TCTTAGTTTTTAAAAAGGCCTTTGGGATTTGCCGGA

**ACGCGT**AAGCGGCCGCGCATCTAGATTCGAAGAAAATGACCGACCAAGCGACGCCCAACCTGCCATCA

CGAGATTTCGATTCCACCGCCGCCTTCTATGAAAGG

**Restriction Sites:** Sgfl-Mlul

OTI Disclaimer: Our molecular clone sequence data has been matched to the sequence identifier above as a

point of reference. Note that the complete sequence of this clone is largely the same as the

reference sequence but may contain minor differences, e.g., single nucleotide

polymorphisms (SNPs).

**Components:** The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The

package also includes 100 pmols of both the corresponding 5' and 3' vector primers in

separate vials.

**RefSeq:** <u>NM 002741.5</u>



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## PKN1 (NM\_002741) Human 3' UTR Clone - SC200626

Summary: The protein encoded by this gene belongs to the protein kinase C superfamily. This kinase is

activated by Rho family of small G proteins and may mediate the Rho-dependent signaling pathway. This kinase can be activated by phospholipids and by limited proteolysis. The 3-phosphoinositide dependent protein kinase-1 (PDPK1/PDK1) is reported to phosphorylate this kinase, which may mediate insulin signals to the actin cytoskeleton. The proteolytic activation of this kinase by caspase-3 or related proteases during apoptosis suggests its role in signal transduction related to apoptosis. Alternatively spliced transcript variants encoding distinct

isoforms have been observed. [provided by RefSeq, Jul 2008]

Locus ID: 5585

MW: 3.7