

Product datasheet for **SC200669**

PSMC3 (NM_002804) Human 3' UTR Clone

Product data:

Product Type:	3' UTR Clones
Product Name:	PSMC3 (NM_002804) Human 3' UTR Clone
Vector:	pMirTarget (PS100062)
Symbol:	PSMC3
Synonyms:	DCIDP; RPT5; TBP1
ACCN:	NM_002804
Insert Size:	90 bp
Insert Sequence:	>SC200669 3'UTR clone of NM_002804 The sequence shown below is from the reference sequence of NM_002804. The complete sequence of this clone may contain minor differences, such as SNPs. Blue=Stop Codon Red=Cloning site GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAAGCCAAGAAGGGCGGAAAGATCGCCGTG TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC AAGAAAGCCAACCTACAATACTACGCCTAGGGCACACAGCCAGCCCCAGTCTCACGGCTGAAGTGCGC AATAAAAGATGGTTTAGGGTC ACGCGTAAGCGGCCGCGCATCTAGATTGAAGAAAATGACCGACCAAGCGACGCCAACCTGCCATCA CGAGATTCGATTCCACCGCCGCTTCTATGAAAGG
Restriction Sites:	SgfI-MluI
OTI Disclaimer:	Our molecular clone sequence data has been matched to the sequence identifier above as a point of reference. Note that the complete sequence of this clone is largely the same as the reference sequence but may contain minor differences , e.g., single nucleotide polymorphisms (SNPs).
Components:	The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The package also includes 100 pmols of both the corresponding 5' and 3' vector primers in separate vials.
RefSeq:	<u>NM_002804.5</u>



[View online »](#)

Summary:

The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes one of the ATPase subunits, a member of the triple-A family of ATPases that have chaperone-like activity. This subunit may compete with PSMC2 for binding to the HIV tat protein to regulate the interaction between the viral protein and the transcription complex. A pseudogene has been identified on chromosome 9. [provided by RefSeq, Jul 2008]

Locus ID:

5702

MW:

3.5