

## Product datasheet for **SC200727**

### Proteasome subunit beta type 4 (PSMB4) (NM\_002796) Human 3' UTR Clone

#### Product data:

Product Type:	3' UTR Clones
Product Name:	Proteasome subunit beta type 4 (PSMB4) (NM_002796) Human 3' UTR Clone
Symbol:	Proteasome subunit beta type 4
Synonyms:	HN3; HsN3; PRAAS3; PROS-26; PROS26
Mammalian Cell Selection:	Neomycin
Vector:	pMirTarget (PS100062)
ACCN:	NM_002796
Insert Size:	138 bp
Insert Sequence:	>SC200727 3'UTR clone of NM_002796 The sequence shown below is from the reference sequence of NM_002796. The complete sequence of this clone may contain minor differences, such as SNPs. <b>Blue</b> =Stop Codon <b>Red</b> =Cloning site  GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAGGCCAAGAAGGGCGGAAAGATCGCCGTG TAACAATTGGCAGAGCTCAGAATTCAA <b>GCGATCGCC</b> ATTGCCACATGATCAGTGGCTTTGAA <b>TGA</b> AATACAGATGCATTATCCAGAAGTGAAGTTGCCCTACTT TTAACTTTGAACTTGGCTAGTTCAAAGATAGACTCTTCTTTGTAAAGTAAATAAATTCTCAAATGC <b>ACGCGT</b> AAGCGGCCGCGCATCTAGATTCAAGAAAATGACCGACCAAGCGACGCCCAACCTGCCATCA CGAGATTCGATTCCACCGCCGCTTCTATGAAAGG
Restriction Sites:	Sgfl-Mlul
OTI Disclaimer:	Our molecular clone sequence data has been matched to the sequence identifier above as a point of reference. Note that the complete sequence of this clone is largely the same as the reference sequence but may contain minor differences , e.g., single nucleotide polymorphisms (SNPs).
Components:	The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The package also includes 100 pmols of both the corresponding 5' and 3' vector primers in separate vials.
RefSeq:	<u><a href="#">NM_002796.3</a></u>



[View online »](#)

**Summary:**

The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the proteasome B-type family, also known as the T1B family, that is a 20S core beta subunit. [provided by RefSeq, Jul 2008]

**Locus ID:**

5692

**MW:**

5