

## **Product datasheet for SC200796**

## RFC4 (NM 181573) Human 3' UTR Clone

## **Product data:**

**Product Type:** 3' UTR Clones

Product Name: RFC4 (NM\_181573) Human 3' UTR Clone

Symbol: RFC4

**Synonyms:** A1; RFC37

Mammalian Cell Neomycin

Selection:

**Vector:** pMirTarget (PS100062)

**ACCN:** NM\_181573

**Insert Size:** 99 bp

Insert Sequence: >SC200796 3'UTR clone of NM\_181573

The sequence shown below is from the reference sequence of NM\_181573. The complete

sequence of this clone may contain minor differences, such as SNPs.

Blue=Stop Codon Red=Cloning site

GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAGGCCAAGAAGGGCGGAAAGATCGCCGTG

TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC

GTGATGCAGCAGTTATCTCAGAATTGTTAACGTGAATATATCTGGATGGGGGGGTTTTGTAAATAATGAA

GTTGTAATAAAAATAAAATGACCAAAAGCA

**ACGCGT**AAGCGGCCGCGCATCTAGATTCGAAGAAAATGACCGACCAAGCGACGCCCAACCTGCCATCA

CGAGATTTCGATTCCACCGCCGCCTTCTATGAAAGG

**Restriction Sites:** Sgfl-Mlul

OTI Disclaimer: Our molecular clone sequence data has been matched to the sequence identifier above as a

point of reference. Note that the complete sequence of this clone is largely the same as the

reference sequence but may contain minor differences, e.g., single nucleotide

polymorphisms (SNPs).

**Components:** The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The

package also includes 100 pmols of both the corresponding 5' and 3' vector primers in

separate vials.

**RefSeq:** NM 181573.3



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



reported. [provided by RefSeq, Jul 2008]

ORIGENE

**Summary:** The elongation of primed DNA templates by DNA polymerase delta and DNA polymerase

epsilon requires the accessory proteins proliferating cell nuclear antigen (PCNA) and replication factor C (RFC). RFC, also named activator 1, is a protein complex consisting of five distinct subunits of 140, 40, 38, 37, and 36 kD. This gene encodes the 37 kD subunit. This subunit forms a core complex with the 36 and 40 kDa subunits. The core complex possesses DNA-dependent ATPase activity, which was found to be stimulated by PCNA in an in vitro system. Alternatively spliced transcript variants encoding the same protein have been

**Locus ID:** 5984

**MW:** 3.9