

Product datasheet for **SC200805**

Caspase 5 (CASP5) (NM_001136109) Human 3' UTR Clone

Product data:

Product Type:	3' UTR Clones
Product Name:	Caspase 5 (CASP5) (NM_001136109) Human 3' UTR Clone
Vector:	pMirTarget (PS100062)
Symbol:	CASP5
Synonyms:	ICE(rel)III; ICEREL-III; ICH-3
ACCN:	NM_001136109
Insert Size:	139 bp
Insert Sequence:	>SC200805 3'UTR clone of NM_001136109 The sequence shown below is from the reference sequence of NM_001136109. The complete sequence of this clone may contain minor differences, such as SNPs. Blue =Stop Codon Red =Cloning site GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAAGCCAAGAAGGGCGGAAAGATCGCCGTG TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC AGAGATTTCTACCTCTTTCTGGCAAT TG AAAATGAAACACAGGCAGCCAGCCCTCTGTCAACA TCAAAGAGCACATTTACCAGTATAGCTTCATAGTCAATATTTGGTATTTCAATAAAAGTAAAGACTGTA ACGCGT AAGCGGCCGCGCATCTAGATT CGA AGAAAATGACCGACCAAGCGACGCCAACCTGCCATCA CGAGATTT CGA TTCACCGCCGCTTCTATGAAAGG
Restriction Sites:	SgfI-MluI
OTI Disclaimer:	Our molecular clone sequence data has been matched to the sequence identifier above as a point of reference. Note that the complete sequence of this clone is largely the same as the reference sequence but may contain minor differences , e.g., single nucleotide polymorphisms (SNPs).
Components:	The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The package also includes 100 pmols of both the corresponding 5' and 3' vector primers in separate vials.
RefSeq:	NM_001136109.3



[View online »](#)

Summary:

This gene encodes a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce two subunits, large and small, that dimerize to form the active enzyme. Overexpression of the active form of this enzyme induces apoptosis in fibroblasts. Max, a central component of the Myc/Max/Mad transcription regulation network important for cell growth, differentiation, and apoptosis, is cleaved by this protein; this process requires Fas-mediated dephosphorylation of Max. The expression of this gene is regulated by interferon-gamma and lipopolysaccharide. Alternatively spliced transcript variants have been identified for this gene. [provided by RefSeq, Aug 2010]

Locus ID:

838

MW:

5.3