

Product datasheet for SC200859

TIMP1 (NM 003254) Human 3' UTR Clone

Product data:

Product Type: 3' UTR Clones

Product Name: TIMP1 (NM 003254) Human 3' UTR Clone

Symbol: TIMP1

Synonyms: CLGI; EPA; EPO; HCI; TIMP; TIMP-1

Mammalian Cell

Selection:

Neomycin

Vector: pMirTarget (PS100062)

ACCN: NM_003254

Insert Size: 128 bp

The sequence shown below is from the reference sequence of NM_003254. The complete

sequence of this clone may contain minor differences, such as SNPs.

Blue=Stop Codon Red=Cloning site

GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAGGCCAAGAAGGGCGGAAAGATCGCCGTG

TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC

TGGCAGTCCCTGCGGTCCCAGATAGCCTGAATCCTGCCCGGAGTGGAAGCTGAAGCCTGCACAGTGTCC

ACCCTGTTCCCACTCCCATCTTTCTTCCGGACAATGAAATAAAGAGTTACCACCCAGCA

ACGCGTAAGCGGCCGCGCATCTAGATTCGAAGAAAATGACCGACCAAGCGACGCCCAACCTGCCATCA

CGAGATTTCGATTCCACCGCCGCCTTCTATGAAAGG

Restriction Sites: Sgfl-Mlul

OTI Disclaimer: Our molecular clone sequence data has been matched to the sequence identifier above as a

point of reference. Note that the complete sequence of this clone is largely the same as the

reference sequence but may contain minor differences, e.g., single nucleotide

polymorphisms (SNPs).

Components: The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The

package also includes 100 pmols of both the corresponding 5' and 3' vector primers in

separate vials.

RefSeq: <u>NM 003254.3</u>



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



ORIGENE

Summary: This gene belongs to the TIMP gene family. The proteins encoded by this gene family are

natural inhibitors of the matrix metalloproteinases (MMPs), a group of peptidases involved in degradation of the extracellular matrix. In addition to its inhibitory role against most of the known MMPs, the encoded protein is able to promote cell proliferation in a wide range of cell types, and may also have an anti-apoptotic function. Transcription of this gene is highly inducible in response to many cytokines and hormones. In addition, the expression from some but not all inactive X chromosomes suggests that this gene inactivation is polymorphic in human females. This gene is located within intron 6 of the synapsin I gene and is

transcribed in the opposite direction. [provided by RefSeq, Jul 2008]

Locus ID: 7076

MW: 4.7