

Product datasheet for SC201110

Caspase 1 (CASP1) (NM_033293) Human 3' UTR Clone

Product data:

Product Type: 3' UTR Clones
Product Name: Caspase 1 (CASP1) (NM_033293) Human 3' UTR Clone
Symbol: Caspase 1
Synonyms: ICE; IL1BC; P45
Mammalian Cell Selection: Neomycin
Vector: pMirTarget (PS100062)
ACCN: NM_033293
Insert Size: 119 bp
Insert Sequence: >SC201110 3' UTR clone of NM_033293
 The sequence shown below is from the reference sequence of NM_033293. The complete sequence of this clone may contain minor differences, such as SNPs. **Red**=Cloning site
Blue=Stop Codon

CAATTGGCAGAGCTCAGAATTCAAGCGATCGC

CTACCTCTTCCCAGGACATTAAATAAGGAACTGTATGAATGTCTGTGGGCAGGAAGTGAAGAGATCCT
 TCTGTAAAGGTTTTTGAATTATGTCTGCTGAATAATAAACTTTTTTGA

ACGCGTAAGCGGCCGCGGCATCTAGATTGAAGAAAATGACCG

Restriction Sites: SgfI-MluI
OTI Disclaimer: Our molecular clone sequence data has been matched to the sequence identifier above as a point of reference. Note that the complete sequence of this clone is largely the same as the reference sequence but may contain minor differences, e.g., single nucleotide polymorphisms (SNPs).
Components: The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The package also includes 100 pmols of both the corresponding 5' and 3' vector primers in separate vials.
RefSeq: NM_033293.2



[View online »](#)

Summary:

This gene encodes a protein which is a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce 2 subunits, large and small, that dimerize to form the active enzyme. This gene was identified by its ability to proteolytically cleave and activate the inactive precursor of interleukin-1, a cytokine involved in the processes such as inflammation, septic shock, and wound healing. This gene has been shown to induce cell apoptosis and may function in various developmental stages. Studies of a similar gene in mouse suggest a role in the pathogenesis of Huntington disease. Alternative splicing results in transcript variants encoding distinct isoforms. [provided by RefSeq, Mar 2012]

Locus ID:

834