

## **Product datasheet for SC201190**

## MRPL53 (NM\_053050) Human 3' UTR Clone

**Product data:** 

Product Type: 3' UTR Clones

Symbol: MRPL53

Synonyms: L53MT

Mammalian Cell Neomycin

Selection:

**Vector:** pMirTarget (PS100062)

**ACCN:** NM\_053050

Insert Size: 167 bp

Insert Sequence: >SC201190 3'UTR clone of NM\_053050

The sequence shown below is from the reference sequence of NM\_053050. The complete sequence of

this clone may contain minor differences, such as SNPs.

Blue=Stop Codon Red=Cloning site

GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAGGCCAAGAAGGGCGGAAAGATCGCCGTG

TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC

GACAAGCCGGGCGCTGATACTGGTCGCTGACAGCGCCCAAAGAGACCAACAAGATGATTTTAGCGTGGAC TAGGACACTTAACCTAAGAAGAGTTTCACTTAATCATTCAAATCACTATCTGAAGGGTCACGGAGCGCA

AAATAAAGTTTAAAACCCTGCTACCACAA

CGAGATTTCGATTCCACCGCCGCCTTCTATGAAAGG

Restriction Sites: Safl-Mlul

Our molecular clone sequence data has been matched to the sequence identifier above as a

point of reference. Note that the complete sequence of this clone is largely the same as the reference sequence but may contain minor differences, e.g., single nucleotide polymorphisms

(SNPs).

Components: The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The

package also includes 100 pmols of both the corresponding 5' and 3' vector primers in

separate vials.



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

EU: info-de@origene.com CN: techsupport@origene.cn



## MRPL53 (NM\_053050) Human 3' UTR Clone | SC201190

Note: Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um

filter is required.

**RefSeq:** <u>NM\_053050.5</u>

Summary: Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in

protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another

difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ

greatly in sequence, and sometimes in biochemical properties, which prevents easy

recognition by sequence homology. This gene encodes a 39S subunit protein. A pseudogene corresponding to this gene is found on chromosome 1p. [provided by RefSeq, Jul 2008]

**Locus ID:** 116540

**MW:** 6.2