

Product datasheet for SC201305

PIG3 (TP53I3) (NM 004881) Human 3' UTR Clone

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com

OriGene Technologies, Inc.

techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

Product data:

Product Type: 3' UTR Clones

Product Name: PIG3 (TP53I3) (NM_004881) Human 3' UTR Clone

Symbol: PIG3 Synonyms: PIG3

Mammalian Cell Neomycin

Selection:

Vector: pMirTarget (PS100062)

ACCN: NM_004881

Insert Size: 176 bp

Insert Sequence: >SC201305 3'UTR clone of NM_004881

The sequence shown below is from the reference sequence of NM_004881. The complete

sequence of this clone may contain minor differences, such as SNPs.

Blue=Stop Codon Red=Cloning site

GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAGGCCAAGAAGGGCGGAAAGATCGCCGTG

TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC

GGCAAGATCGTCCTGGAACTGCCCCAGTGAAGGAGGATGGGGCAGGACAGGACGCGGCCACCCCAGGCC
TTTCCAGAGCAAACCTGGAGAAGATTCACAATAGACAGGCCAAGAAACCCGGTGCTTCCTCCAGAGCCG

TTTAAAGCTGATATGAGGAAATAAAGAGTGAACTGGAA

CGAGATTTCGATTCCACCGCCGCCTTCTATGAAAGG

Restriction Sites: Sgfl-Mlul

OTI Disclaimer: Our molecular clone sequence data has been matched to the sequence identifier above as a

point of reference. Note that the complete sequence of this clone is largely the same as the

reference sequence but may contain minor differences, e.g., single nucleotide

polymorphisms (SNPs).

Components: The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The

package also includes 100 pmols of both the corresponding 5' and 3' vector primers in

separate vials.

RefSeq: <u>NM 004881.5</u>





ORIGENE

Summary:

MW:

The protein encoded by this gene is similar to oxidoreductases, which are enzymes involved in cellular responses to oxidative stresses and irradiation. This gene is induced by the tumor suppressor p53 and is thought to be involved in p53-mediated cell death. It contains a p53 consensus binding site in its promoter region and a downstream pentanucleotide microsatellite sequence. P53 has been shown to transcriptionally activate this gene by interacting with the downstream pentanucleotide microsatellite sequence. The microsatellite is polymorphic, with a varying number of pentanucleotide repeats directly correlated with the extent of transcriptional activation by p53. It has been suggested that the microsatellite polymorphism may be associated with differential susceptibility to cancer. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, May 2011]

Locus ID: 9540 7