

Product datasheet for SC201313

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Carboxypeptidase Z (CPZ) (NM_001014447) Human 3' UTR Clone

Product data:

Product Type: 3' UTR Clones

Symbol: Carboxypeptidase Z

Mammalian Cell Neomycin

Selection:

Vector: pMirTarget (PS100062)

ACCN: NM_001014447

Insert Size: 165 bp

Insert Sequence: >SC201313 3'UTR clone of NM_001014447

The sequence shown below is from the reference sequence of NM_001014447. The complete sequence

of this clone may contain minor differences, such as SNPs.

Blue=Stop Codon Red=Cloning site

GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAGGCCAAGAAGGGCGGAAAGATCGCCGTG

TAACAATTGGCAGAGCTCAGAATTCAA<mark>GCGATCGC</mark>C

CACAGGCCACGCTGGCTCAAGTACTAGCCCCGGCCCCAGCACCCCAGGATGTGGAGACCGAGGCCCCATCTCCGCATCCCGGGCTCCTGGCTCTTGATTTTGTCTGCCACAGACATCCCACAAAGCCGCTGCCA

TTTTATTAAAGTGTTTTGATCCACTTT

ACGCGTAAGCGGCCGCGCATCTAGATTCGAAGAAAATGACCGACCAAGCGACGCCCAACCTGCCATCA

CGAGATTTCGATTCCACCGCCGCCTTCTATGAAAGG

Restriction Sites: Sgfl-Mlul

OTI Disclaimer: Our molecular clone sequence data has been matched to the sequence identifier above as a

point of reference. Note that the complete sequence of this clone is largely the same as the reference sequence but may contain minor differences, e.g., single nucleotide polymorphisms

(SNPs).

Components: The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The

package also includes 100 pmols of both the corresponding 5' and 3' vector primers in

separate vials.

Note: Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um

filter is required.





Carboxypeptidase Z (CPZ) (NM_001014447) Human 3' UTR Clone | SC201313

RefSeq: <u>NM_001014447.3</u>

Summary: This gene encodes a member of the metallocarboxypeptidase family. This enzyme displays

carboxypeptidase activity towards substrates with basic C-terminal residues. It is most active at neutral pH and is inhibited by active site-directed inhibitors of metallocarboxypeptidases. Alternative splicing in the coding region results in multiple transcript variants encoding

different isoforms. [provided by RefSeq, Jul 2008]

Locus ID: 8532

MW: 6.4