

## **Product datasheet for SC201319**

## DHPS (NM 001930) Human 3' UTR Clone

**Product data:** 

**Product Type:** 3' UTR Clones

Product Name: DHPS (NM\_001930) Human 3' UTR Clone

**Vector:** pMirTarget (PS100062)

Symbol: DHPS

Synonyms: DHS; DS; MIG13; NEDSSWI

**ACCN:** NM\_001930

**Insert Size:** 151 bp

Insert Sequence: >SC201319 3'UTR clone of NM\_001930

The sequence shown below is from the reference sequence of NM\_001930. The complete

sequence of this clone may contain minor differences, such as SNPs.

Blue=Stop Codon Red=Cloning site

GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAGGCCAAGAAGGGCGGAAAGATCGCCGTG

TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC

TGAAGTCCTTCCA

**ACGCGT**AAGCGGCCGCGCATCTAGATTCGAAGAAAATGACCGACCAAGCGACGCCCAACCTGCCATCA

CGAGATTTCGATTCCACCGCCGCCTTCTATGAAAGG

**Restriction Sites:** Sgfl-Mlul

OTI Disclaimer: Our molecular clone sequence data has been matched to the sequence identifier above as a

point of reference. Note that the complete sequence of this clone is largely the same as the

reference sequence but may contain minor differences, e.g., single nucleotide

polymorphisms (SNPs).

**Components:** The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The

package also includes 100 pmols of both the corresponding 5' and 3' vector primers in

separate vials.

**RefSeg:** NM 001930.4



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## DHPS (NM\_001930) Human 3' UTR Clone - SC201319

**Summary:** This gene encodes a protein that is required for the formation of hypusine, a unique amino

acid formed by the posttranslational modification of only one protein, eukaryotic translation initiation factor 5A. The encoded protein catalyzes the first step in hypusine formation by transferring the butylamine moiety of spermidine to a specific lysine residue of the eukaryotic translation initiation factor 5A precursor, forming an intermediate deoxyhypusine residue. Alternatively spliced transcript variants encoding multiple isoforms have been observed for

this gene. [provided by RefSeq, May 2011]

**Locus ID:** 1725

**MW:** 5.5