

Product datasheet for SC201380

TROY (TNFRSF19) (NM_018647) Human 3' UTR Clone

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	3' UTR Clones
Product Name:	TROY (TNFRSF19) (NM_018647) Human 3' UTR Clone
Symbol:	TROY
Synonyms:	TAJ; TAJ-alpha; TRADE; TROY
Mammalian Cell Selection:	Neomycin
Vector:	pMirTarget (PS100062)
ACCN:	NM_018647
Insert Size:	177 bp
Insert Sequence:	<pre>>SC201380 3'UTR clone of NM_018647 The sequence shown below is from the reference sequence of NM_018647. The complete sequence of this clone may contain minor differences, such as SNPs. Blue=Stop Codon Red=Cloning site GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAGGCCAAGAAGGGCGGAAAGATCGCCGTG TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC CAGGTAAGGCAGCGACTGGGTTCCCTGTGAACACAGCACTGACTTACAGTAGATCAGAACTCTGTTCCC AGCATAAGATTTGGGGGAACCTGATGAGTTTTTTTTTGCATCTTAATAATTTCTTGTATGTTGTAGA GTATGTTTTAAAATAAATTTCAAGTATTTTTTTAAAAAA ACGCGTAAGCGGCCGCGGCATCTAGATTCGAAGAAAATGACCGACC</pre>
Restriction Sites:	SgfI-Mlul
OTI Disclaimer:	Our molecular clone sequence data has been matched to the sequence identifier above as a point of reference. Note that the complete sequence of this clone is largely the same as the reference sequence but may contain minor differences , e.g., single nucleotide polymorphisms (SNPs).
Components:	The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The package also includes 100 pmols of both the corresponding 5' and 3' vector primers in separate vials.
RefSeq:	<u>NM 018647.5</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	TROY (TNFRSF19) (NM_018647) Human 3' UTR Clone – SC201380
Summary:	The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is highly expressed during embryonic development. It has been shown to interact with TRAF family members, and to activate JNK signaling pathway when overexpressed in cells. This receptor is capable of inducing apoptosis by a caspase-independent mechanism, and it is thought to play an essential role in embryonic development. Alternatively spliced transcript variants encoding distinct isoforms have been described. [provided by RefSeq, Jul 2008]
Locus ID:	55504
MW:	7

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US