

## **Product datasheet for SC201406**

## RFXANK (NM 134440) Human 3' UTR Clone

**Product data:** 

**Product Type:** 3' UTR Clones

Product Name: RFXANK (NM\_134440) Human 3' UTR Clone

Symbol: RFXANK

Synonyms: ANKRA1; BLS; F14150\_1; RFX-B

Mammalian Cell

Selection:

Neomycin

**Vector:** pMirTarget (PS100062)

**ACCN:** NM\_134440

**Insert Size:** 177 bp

The sequence shown below is from the reference sequence of NM\_134440. The complete

sequence of this clone may contain minor differences, such as SNPs.

Blue=Stop Codon Red=Cloning site

GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAGGCCAAGAAGGGCGGAAAGATCGCCGTG

TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC

AGCAACCTGGTGCCCGCTGACCCTGAGTGAAGGCCGCCTGCCGGGGACTCAGACACTCAGGGAACAAAA TGGTCAGCCAGAGCTGGGGAAACCCAGAACTGACTTCAAAGGCAGCTTCTGGACAGGTGGTGGGAGGGG

ACCCTTCCCAAGAGGAACCAATAAACCTTCTGTGCAGAA

CGAGATTTCGATTCCACCGCCGCCTTCTATGAAAGG

**Restriction Sites:** Sgfl-Mlul

OTI Disclaimer: Our molecular clone sequence data has been matched to the sequence identifier above as a

point of reference. Note that the complete sequence of this clone is largely the same as the

reference sequence but may contain minor differences, e.g., single nucleotide

polymorphisms (SNPs).

**Components:** The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The

package also includes 100 pmols of both the corresponding 5' and 3' vector primers in

separate vials.

**RefSeq:** <u>NM 134440.3</u>



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## RFXANK (NM\_134440) Human 3' UTR Clone - SC201406

**Summary:** 

Major histocompatibility (MHC) class II molecules are transmembrane proteins that have a central role in development and control of the immune system. The protein encoded by this gene, along with regulatory factor X-associated protein and regulatory factor-5, forms a complex that binds to the X box motif of certain MHC class II gene promoters and activates their transcription. Once bound to the promoter, this complex associates with the non-DNA-binding factor MHC class II transactivator, which controls the cell type specificity and inducibility of MHC class II gene expression. This protein contains ankyrin repeats involved in protein-protein interactions. Mutations in this gene have been linked to bare lymphocyte syndrome type II, complementation group B. Multiple alternatively spliced transcript variants encoding different isoforms have been described for this gene. [provided by RefSeq, Jul 2013]

**Locus ID:** 8625 **MW:** 6.7