

## Product datasheet for **SC201425**

### **MCM7 (NM\_182776) Human 3' UTR Clone**

#### **Product data:**

|                    |   |
|--------------------|---|
| Product Type:      | 3' UTR Clones   |
| Product Name:      | MCM7 (NM_182776) Human 3' UTR Clone   |
| Vector:            | pMirTarget (PS100062)   |
| Symbol:            | MCM7  |
| Synonyms:          | CDC47; MCM2; P1.1-MCM3; P1CDC47; P85MCM; PNAS146; PPP1R104  |
| ACCN:              | NM_182776   |
| Insert Size:       | 234 bp  |
| Insert Sequence:   | >SC201425 3'UTR clone of NM_182776<br>The sequence shown below is from the reference sequence of NM_182776. The complete sequence of this clone may contain minor differences, such as SNPs.<br><b>Blue</b> =Stop Codon <b>Red</b> =Cloning site<br><br>GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAAGCCAAGAAGGGCGGAAAGATCGCCGTG<br>TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC<br>GCTTCCCGGACACGGATCACTTTTGTCTGATTCCAGCCTGCTTGCAACCCTGGGGTCTCTGTTCCT<br>GCTGGCCTGCCCTTGGGAAGGGGAGGAGGAGCCCTCTTTCTCCCATG<br>CTGCCTTACTCCTTTTGCTAATAAAAAGTGTGTAGATTGTCATCTTCTAGCCTGGGCCTGACTTCCA<br>TTAAAACAGGGTTTTGTGCGTTTTTTA<br><b>ACGCGT</b> AAGCGGCCGCGCATCTAGATTGAAAGAAAATGACCGACCAAGCGACGCCCAACCTGCCATCA<br>CGAGATTCGATTCCACCGCCCTTCTATGAAAGG |
| Restriction Sites: | SgfI-MluI   |
| OTI Disclaimer:    | Our molecular clone sequence data has been matched to the sequence identifier above as a point of reference. Note that the complete sequence of this clone is largely the same as the reference sequence but may contain minor differences, e.g., single nucleotide polymorphisms (SNPs).   |
| Components:        | The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The package also includes 100 pmols of both the corresponding 5' and 3' vector primers in separate vials.  |
| RefSeq:            | <u><a href="#">NM_182776.3</a></u>  |



[View online »](#)

**Summary:**

The protein encoded by this gene is one of the highly conserved mini-chromosome maintenance proteins (MCM) that are essential for the initiation of eukaryotic genome replication. The hexameric protein complex formed by the MCM proteins is a key component of the pre-replication complex (pre\_RC) and may be involved in the formation of replication forks and in the recruitment of other DNA replication related proteins. The MCM complex consisting of this protein and MCM2, 4 and 6 proteins possesses DNA helicase activity, and may act as a DNA unwinding enzyme. Cyclin D1-dependent kinase, CDK4, is found to associate with this protein, and may regulate the binding of this protein with the tumorsuppressor protein RB1/RB. Alternatively spliced transcript variants encoding distinct isoforms have been reported. [provided by RefSeq, Jul 2008]

**Locus ID:**

4176

**MW:**

8.5