

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for SC201547

Macrophage inflammatory protein 5 (CCL15) (NM_032965) Human 3' UTR Clone

Product data:

Product Type:	3' UTR Clones
Product Name:	Macrophage inflammatory protein 5 (CCL15) (NM_032965) Human 3' UTR Clone
Vector:	pMirTarget (PS100062)
Symbol:	CCL15
Synonyms:	HCC-2; HMRP-2B; LKN-1; LKN1; MIP-1 delta; MIP-1D; MIP-5; MRP-2B; NCC-3; NCC3; SCYA15; SCYL3; SY15
ACCN:	NM_032965
Insert Size:	179 bp
Insert Sequence:	>SC201547 3' UTR clone of NM_032965 The sequence shown below is from the reference sequence of NM_032965. The complete sequence of this clone may contain minor differences, such as SNPs. Red=Cloning site Blue=Stop Codon
	CAATTGGCAGAGCTCAGAATTCAAGCGATCGC
	CGGGAGTTCAGGATTGCATGAAAAAGCTGAAGCCCTACTCAATA TAA TAATAAAGAGACAAAAGAGGCCA GCCACCCACCTCCAACACCTCCTGTGAGTTTCTTGGTCTGAAATACTTAAAAAAATATATAT
	ACGCGTAAGCGGCCGCGCATCTAGATTCGAAGAAAATGACCG
Restriction Sites:	Sgfl-Mlul
OTI Disclaimer:	Our molecular clone sequence data has been matched to the sequence identifier above as a point of reference. Note that the complete sequence of this clone is largely the same as the reference sequence but may contain minor differences , e.g., single nucleotide polymorphisms (SNPs).
Components:	The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The package also includes 100 pmols of both the corresponding 5' and 3' vector primers in separate vials.
RefSeq:	<u>NM 032965.3</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	Macrophage inflammatory protein 5 (CCL15) (NM_032965) Human 3' UTR Clone – SC201547
Summary:	This gene is located in a cluster of similar genes in the same region of chromosome 17. These genes encode CC cytokines, which are secreted proteins characterized by two adjacent cysteines. The product of this gene is chemotactic for T cells and monocytes, and acts through C-C chemokine receptor type 1 (CCR1). The proprotein is further processed into numerous smaller functional peptides. Naturally-occurring readthrough transcripts occur from this gene into the downstream gene, CCL14 (chemokine (C-C motif) ligand 14). [provided by RefSeq, Jan 2013]
Locus ID:	6359

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US