

## Product datasheet for **SC201629**

### Sec8 (EXOC4) (NM\_001037126) Human 3' UTR Clone

#### Product data:

Product Type:	3' UTR Clones
Product Name:	Sec8 (EXOC4) (NM_001037126) Human 3' UTR Clone
Symbol:	Sec8
Synonyms:	SEC8; SEC8L1; Sec8p
Mammalian Cell Selection:	Neomycin
Vector:	pMirTarget (PS100062)
ACCN:	NM_001037126
Insert Size:	183 bp
Insert Sequence:	>SC201629 3'UTR clone of NM_001037126 The sequence shown below is from the reference sequence of NM_001037126. The complete sequence of this clone may contain minor differences, such as SNPs. <b>Blue</b> =Stop Codon <b>Red</b> =Cloning site  GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAAGCCAAGAAGGGCGGAAAGATCGCCGTG TAACAATTGGCAGAGCTCAGAATTCAA <b>GCGATCGCC</b> TATAGTCGGAGTGGAGAAGTCAAGGG <b>TAG</b> CGGCGAAGAAATCCACCAGAAAAGACAATCAAAGTAACAT TAAAAAGCTCAAATTTTACAAAAGTTAAGTTCTGCCAACATTTTCAATTTATCATACAATTAGAAATG TGACTCAGTCTAACAAACCCCAAGAGGATAAAAAGGATTA <b>ACGCGT</b> AAGCGGCCGCGCATCTAGATTGAAGAAAATGACCGACCAAGCGACGCCCAACCTGCCATCA CGAGATTTGATTCCACCGCCGCTTCTATGAAAGG
Restriction Sites:	Sgfl-MluI
OTI Disclaimer:	Our molecular clone sequence data has been matched to the sequence identifier above as a point of reference. Note that the complete sequence of this clone is largely the same as the reference sequence but may contain minor differences, e.g., single nucleotide polymorphisms (SNPs).
Components:	The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The package also includes 100 pmols of both the corresponding 5' and 3' vector primers in separate vials.
RefSeq:	<a href="#">NM_001037126.2</a>



[View online »](#)

**Summary:**

The protein encoded by this gene is a component of the exocyst complex, a multiple protein complex essential for targeting exocytic vesicles to specific docking sites on the plasma membrane. Though best characterized in yeast, the component proteins and functions of exocyst complex have been demonstrated to be highly conserved in higher eukaryotes. At least eight components of the exocyst complex, including this protein, are found to interact with the actin cytoskeletal remodeling and vesicle transport machinery. The complex is also essential for the biogenesis of epithelial cell surface polarity. Alternate transcriptional splice variants, encoding different isoforms, have been characterized. [provided by RefSeq, Jul 2008]

**Locus ID:**

60412

**MW:**

7.2