

## Product datasheet for SC201713

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

## ICAM2 (NM 001099787) Human 3' UTR Clone

**Product data:** 

**Product Type:** 3' UTR Clones

**Product Name:** ICAM2 (NM 001099787) Human 3' UTR Clone

Vector: pMirTarget (PS100062)

Symbol: ICAM2 Synonyms: CD102

ACCN: NM 001099787

Insert Size: 182 bp

>SC201713 3'UTR clone of NM\_001099787 **Insert Sequence:** 

The sequence shown below is from the reference sequence of NM\_001099787. The complete

sequence of this clone may contain minor differences, such as SNPs.

Blue=Stop Codon Red=Cloning site

GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAGGCCAAGAAGGGCGGAAAGATCGCCGTG

TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC

AGGAGGCTGCCCCAGGCCTTCCGGCCATAGCAACCATGAGTGGCATGGCCACCACCACGGTGGTCACTG GAACTCAGTGTGACTCCTCAGGGTTGAGGTCCAGCCCTGGCTGAAGGACTGTGACAGGCAGCAGAGACT

TGGGACATTGCCTTTTCTAGCCCGAATACAAACACCTGGACTTA

CGAGATTTCGATTCCACCGCCGCCTTCTATGAAAGG

**Restriction Sites:** Sgfl-Mlul

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the sequence identifier above as a

point of reference. Note that the complete sequence of this clone is largely the same as the

reference sequence but may contain minor differences, e.g., single nucleotide

polymorphisms (SNPs).

Components: The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The

package also includes 100 pmols of both the corresponding 5' and 3' vector primers in

separate vials.

RefSeq: NM 001099787.2





## ICAM2 (NM\_001099787) Human 3' UTR Clone - SC201713

Summary: The protein encoded by this gene is a member of the intercellular adhesion molecule (ICAM)

family. All ICAM proteins are type I transmembrane glycoproteins, contain 2-9

immunoglobulin-like C2-type domains, and bind to the leukocyte adhesion LFA-1 protein. This

protein may play a role in lymphocyte recirculation by blocking LFA-1-dependent cell

adhesion. It mediates adhesive interactions important for antigen-specific immune response,

NK-cell mediated clearance, lymphocyte recirculation, and other cellular interactions

important for immune response and surveillance. Several transcript variants encoding the

same protein have been found for this gene. [provided by RefSeq, Jul 2008]

**Locus ID:** 3384

**MW:** 6.7