

## Product datasheet for SC201801

## MTH1 (NUDT1) (NM 198949) Human 3' UTR Clone

**Product data:** 

**Product Type:** 3' UTR Clones

**Product Name:** MTH1 (NUDT1) (NM 198949) Human 3' UTR Clone

Symbol: MTH1 Synonyms:

**Mammalian Cell** Neomycin

Selection:

Vector:

pMirTarget (PS100062)

ACCN: NM 198949

**Insert Size:** 174 bp

**Insert Sequence:** >SC201801 3'UTR clone of NM\_198949

The sequence shown below is from the reference sequence of NM\_198949. The complete

sequence of this clone may contain minor differences, such as SNPs.

Blue=Stop Codon Red=Cloning site

GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAGGCCAAGAAGGGCGGAAAGATCGCCGTG

TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC

TACACACTCCGCGAGGTGGACACGGTCTAGCGGGAGCCCCAGGGCAGCCCCTGGGCAGGAGACGTGGCTG CTGAACAGCCGCAAACCATCTTCACCTGGGGGCATTGAGTGGCGCAGAGCCGGGTTTCATCTGGAATTA

ACTGGATGGAAGGGAAAATAAAGCTATCTAGCGGTG

CGAGATTTCGATTCCACCGCCGCCTTCTATGAAAGG

**Restriction Sites:** Sgfl-Mlul

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the sequence identifier above as a

point of reference. Note that the complete sequence of this clone is largely the same as the

reference sequence but may contain minor differences, e.g., single nucleotide

polymorphisms (SNPs).

The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The Components:

package also includes 100 pmols of both the corresponding 5' and 3' vector primers in

separate vials.

RefSeq: NM 198949.2



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com





**Summary:** 

Misincorporation of oxidized nucleoside triphosphates into DNA/RNA during replication and transcription can cause mutations that may result in carcinogenesis or neurodegeneration. The protein encoded by this gene is an enzyme that hydrolyzes oxidized purine nucleoside triphosphates, such as 8-oxo-dGTP, 8-oxo-dATP, 2-hydroxy-dATP, and 2-hydroxy rATP, to monophosphates, thereby preventing misincorporation. The encoded protein is localized mainly in the cytoplasm, with some in the mitochondria, suggesting that it is involved in the sanitization of nucleotide pools both for nuclear and mitochondrial genomes. Several alternatively spliced transcript variants, some of which encode distinct isoforms, have been identified. Additional variants have been observed, but their full-length natures have not been determined. A rare single-nucleotide polymorphism that results in the production of an additional, longer isoform (p26) has been described. [provided by RefSeq, Dec 2018]

**Locus ID:** 4521 **MW:** 6.6