

## Product datasheet for **SC201947**

### **ATP5MF (NM\_004889) Human 3' UTR Clone**

#### **Product data:**

<b>Product Type:</b>	3' UTR Clones
<b>Product Name:</b>	ATP5MF (NM_004889) Human 3' UTR Clone
<b>Vector:</b>	pMirTarget (PS100062)
<b>Symbol:</b>	ATP5MF
<b>Synonyms:</b>	ATP5J2; ATP5JL
<b>ACCN:</b>	NM_004889
<b>Insert Size:</b>	162 bp
<b>Insert Sequence:</b>	>SC201947 3'UTR clone of NM_004889 The sequence shown below is from the reference sequence of NM_004889. The complete sequence of this clone may contain minor differences, such as SNPs. <b>Blue</b> =Stop Codon <b>Red</b> =Cloning site  GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAAGGCCAAGAAGGGCGGAAAGATCGCCGTG TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC AAGCACGAGCGGCTCCGCAAATACCACTGAGAGGACACACTCTGCACCCCCACCCCACGACCTTGG CCCGAGCCCCTCCGTGAGGAACACAATCTCAATCGTTGCTGAATCCTTTCATATCCTAATAGGAATTAA CCTCCAAATAAAACATGACTGGTA <b>ACGCGT</b> AAGCGGCCGCGCATCTAGATTGAAAGAAAATGACCGACCAAGCGACGCCCAACCTGCCATCA CGAGATTCGATTCCACCGCCGCTTCTATGAAAGG
<b>Restriction Sites:</b>	Sgfl-MluI
<b>OTI Disclaimer:</b>	Our molecular clone sequence data has been matched to the sequence identifier above as a point of reference. Note that the complete sequence of this clone is largely the same as the reference sequence but may contain minor differences , e.g., single nucleotide polymorphisms (SNPs).
<b>Components:</b>	The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The package also includes 100 pmols of both the corresponding 5' and 3' vector primers in separate vials.
<b>RefSeq:</b>	<u><a href="#">NM_004889.5</a></u>



[View online »](#)

**Summary:**

Mitochondrial ATP synthase catalyzes ATP synthesis, utilizing an electrochemical gradient of protons across the inner membrane during oxidative phosphorylation. It is composed of two linked multi-subunit complexes: the soluble catalytic core, F1, and the membrane-spanning component, Fo, which comprises the proton channel. The catalytic portion of mitochondrial ATP synthase consists of five different subunits (alpha, beta, gamma, delta, and epsilon) assembled with a stoichiometry of 3 alpha, 3 beta, and single representatives of the gamma, delta, and epsilon subunits. The proton channel likely has nine subunits (a, b, c, d, e, f, g, F6 and 8). This gene encodes the f subunit of the Fo complex. Alternatively spliced transcript variants encoding different isoforms have been identified for this gene. This gene has multiple pseudogenes. Naturally occurring read-through transcription also exists between this gene and the downstream pentatricopeptide repeat domain 1 (PTCD1) gene. [provided by RefSeq, Nov 2010]

**Locus ID:** 9551

**MW:** 6.6