

## Product datasheet for SC202118

### Peroxiredoxin 5 (PRDX5) (NM\_181652) Human 3' UTR Clone

#### Product data:

Product Type:	3' UTR Clones
Product Name:	Peroxiredoxin 5 (PRDX5) (NM_181652) Human 3' UTR Clone
Symbol:	Peroxiredoxin 5
Synonyms:	ACR1; AOEB166; B166; HEL-S-55; PLP; PMP20; PRDX6; prx-V; PRXV; SBBI10
Mammalian Cell Selection:	Neomycin
Vector:	pMirTarget (PS100062)
ACCN:	NM_181652
Insert Size:	150 bp
Insert Sequence:	<p>&gt;SC202118 3'UTR clone of NM_181652</p> <p>The sequence shown below is from the reference sequence of NM_181652. The complete sequence of this clone may contain minor differences, such as SNPs.</p> <p>Blue=Stop Codon Red=Cloning site</p>

```

GGCAAGTTGGACGCCGCAAGATCCGCGAGATTCTCATTAAGGCCAAGAAGGCGGAAAGATCGCCGTG
TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC
CTGGCACCCAATATCATCTCACAGCTCTGAGGCCCTGGGCCAGATTACTTCTCCACCCCTCCCTATCT
CACCTGCCAGCCCTGTGCTGGGGCCCTGCAATTGGAATGTTGGCCAGATTCTGCAATAAACACTTGT
GGTTTGGGCCA
ACGCGTAAGCGGCCGCGCATCTAGATTGGAAGAAATGACCGACCAAGCGACGCCAACCTGCCATCA
CGAGATTCGATTCCACCGCCGCTTCTATGAAAGG
  
```

Restriction Sites:	Sgfl-MluI
OTI Disclaimer:	Our molecular clone sequence data has been matched to the sequence identifier above as a point of reference. Note that the complete sequence of this clone is largely the same as the reference sequence but may contain minor differences , e.g., single nucleotide polymorphisms (SNPs).
Components:	The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The package also includes 100 pmols of both the corresponding 5' and 3' vector primers in separate vials.
RefSeq:	<u><a href="#">NM_181652.3</a></u>


[View online »](#)

<b>Summary:</b>	This gene encodes a member of the peroxiredoxin family of antioxidant enzymes, which reduce hydrogen peroxide and alkyl hydroperoxides. The encoded protein interacts with peroxisome receptor 1 and plays an antioxidant protective role in different tissues under normal conditions and during inflammatory processes. The use of alternate transcription start sites is thought to result in transcript variants that use different in-frame translational start codons to generate isoforms that are targeted to the mitochondrion (isoform L) or peroxisome/cytoplasm (isoform S). Multiple related pseudogenes have been defined for this gene. [provided by RefSeq, Nov 2017]
<b>Locus ID:</b>	25824
<b>MW:</b>	5.3