

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for SC202197

Ribonuclease Inhibitor (RNH1) (NM_203388) Human 3' UTR Clone

Product data:

Product Type:	3' UTR Clones
Product Name:	Ribonuclease Inhibitor (RNH1) (NM_203388) Human 3' UTR Clone
Vector:	pMirTarget (PS100062)
Symbol:	RNH1
Synonyms:	RAI; RNH
ACCN:	NM_203388
Insert Size:	206 bp
Insert Sequence:	<pre>>SC202197 3'UTR clone of NM_203388 The sequence shown below is from the reference sequence of NM_203388. The complete sequence of this clone may contain minor differences, such as SNPs. Blue=Stop Codon Red=Cloning site GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAGGCCAAGAAGGGCGGAAAGATCGCCGTG TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC GACAAGCCATCCCTGAGGGTCATCTCCTGAGGCTCTTCCTGCTGCTGCTCTCCCTGGACGACCGGCCTC GAGGCAACCCTGGGGCCCACCAGCCCTGCCATGCTCTCACCCTGCATATCCTAGGTTTGAAGAGAAAC GCTCAGATCCGCTTATTTCTGCCAGTATATTTTGGACACTTTATAATCATTAAAGCACTTTCTTGGCA ACGCGTAAGCGGCCGCGGCATCTAGATTCGAAGAAAATGACCGACC</pre>
Restriction Sites:	Sgfl-Mlul
OTI Disclaimer:	Our molecular clone sequence data has been matched to the sequence identifier above as a point of reference. Note that the complete sequence of this clone is largely the same as the reference sequence but may contain minor differences , e.g., single nucleotide polymorphisms (SNPs).
Components:	The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The package also includes 100 pmols of both the corresponding 5' and 3' vector primers in separate vials.
RefSeq:	<u>NM 203388.3</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	Ribonuclease Inhibitor (RNH1) (NM_203388) Human 3' UTR Clone – SC202197
Summary:	Placental ribonuclease inhibitor (PRI) is a member of a family of proteinaceous cytoplasmic RNase inhibitors that occur in many tissues and bind to both intracellular and extracellular RNases (summarized by Lee et al., 1988 [PubMed 3219362]). In addition to control of intracellular RNases, the inhibitor may have a role in the regulation of angiogenin (MIM 105850). Ribonuclease inhibitor, of 50,000 Da, binds to ribonucleases and holds them in a latent form. Since neutral and alkaline ribonucleases probably play a critical role in the turnover of RNA in eukaryotic cells, RNH may be essential for control of mRNA turnover; the interaction of eukaryotic cells with ribonuclease may be reversible in vivo.[supplied by OMIM, Jul 2010]
Locus ID:	6050
MW:	7.5

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US