

Product datasheet for **SC202280**

DAXX (NM_001141970) Human 3' UTR Clone

Product data:

Product Type:	3' UTR Clones
Product Name:	DAXX (NM_001141970) Human 3' UTR Clone
Vector:	pMirTarget (PS100062)
Symbol:	DAXX
Synonyms:	BING2; DAP6; EAP1; SMIM40
ACCN:	NM_001141970
Insert Size:	215 bp
Insert Sequence:	>SC202280 3'UTR clone of NM_001141970 The sequence shown below is from the reference sequence of NM_001141970. The complete sequence of this clone may contain minor differences, such as SNPs. Blue =Stop Codon Red =Cloning site GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAAGCCAAGAAGGGCGGAAAGATCGCCGTG TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC GAGATCATCGTGCTCTCAGACTCTGATTAGCTGCCTCCCCTTCTCCCTGCCTCCAGAATGTTCTGGGAT AACATTTGGAGGAAGGTGGGAAGCAGATGACTGAGGAAGGGATGGACTAAGCTAATCCCCTTTTGGTGG TGTTTCTTTAAAAAAGCTTAAGTTTTACACAGAAACATTAATAACAATAAAGTTCTTTTC TTACTGTA ACGCGT AAGCGGCCGCGGCATCTAGATTGGAAGAAATGACCGACCAAGCGACGCCCAACCTGCCATCA CGAGATTCGATTCCACCGCCGCTTCTATGAAAGG
Restriction Sites:	SgfI-MluI
OTI Disclaimer:	Our molecular clone sequence data has been matched to the sequence identifier above as a point of reference. Note that the complete sequence of this clone is largely the same as the reference sequence but may contain minor differences, e.g., single nucleotide polymorphisms (SNPs).
Components:	The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The package also includes 100 pmols of both the corresponding 5' and 3' vector primers in separate vials.
RefSeq:	<u>NM_001141970.2</u>



[View online »](#)

Summary:

This gene encodes a multifunctional protein that resides in multiple locations in the nucleus and in the cytoplasm. It interacts with a wide variety of proteins, such as apoptosis antigen Fas, centromere protein C, and transcription factor erythroblastosis virus E26 oncogene homolog 1. In the nucleus, the encoded protein functions as a potent transcription repressor that binds to sumoylated transcription factors. Its repression can be relieved by the sequestration of this protein into promyelocytic leukemia nuclear bodies or nucleoli. This protein also associates with centromeres in G2 phase. In the cytoplasm, the encoded protein may function to regulate apoptosis. The subcellular localization and function of this protein are modulated by post-translational modifications, including sumoylation, phosphorylation and polyubiquitination. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Nov 2008]

Locus ID:

1616

MW:

8.7