

Product datasheet for **SC202292**

UBE2C (NM_007019) Human 3' UTR Clone

Product data:

Product Type:	3' UTR Clones
Product Name:	UBE2C (NM_007019) Human 3' UTR Clone
Vector:	pMirTarget (PS100062)
Symbol:	UBE2C
Synonyms:	dj447F3.2; UBCH10
ACCN:	NM_007019
Insert Size:	215 bp
Insert Sequence:	>SC202292 3'UTR clone of NM_007019 The sequence shown below is from the reference sequence of NM_007019. The complete sequence of this clone may contain minor differences, such as SNPs. Blue =Stop Codon Red =Cloning site GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAAGCCAAGAAGGGCGGAAAGATCGCCGTG TAACAATTGGCAGAGCTCAGAATTCAA GCGATCGCC TCAAAGCAGGTCACCAGCCAGGAGCC TGA CCCAGGCTGCCAGCCTGTCCTTGTGTGCTCTTTTTAAT TTTTCTTAGATGGTCTGTCCTTTTTGTGATTTCTGTATAGGACTCTTTATCTTGAGCTGTGGTATTTT TGTTTTGTTTTGCTTTTTAAATTAAGCCTCGGTTGAGCCCTGTATATTAATAAATGCATTTTTGTC CTTTTTTA ACGCGT AAGCGGCCGCGCATCTAGATTCTGAAGAAAATGACCGACCAAGCGACGCCCAACCTGCCATCA CGAGATTCGATTCCACCGCCGCTTCTATGAAAGG
Restriction Sites:	SgfI-MluI
OTI Disclaimer:	Our molecular clone sequence data has been matched to the sequence identifier above as a point of reference. Note that the complete sequence of this clone is largely the same as the reference sequence but may contain minor differences, e.g., single nucleotide polymorphisms (SNPs).
Components:	The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The package also includes 100 pmols of both the corresponding 5' and 3' vector primers in separate vials.
RefSeq:	<u>NM_007019.4</u>



[View online »](#)

Summary:

The modification of proteins with ubiquitin is an important cellular mechanism for targeting abnormal or short-lived proteins for degradation. Ubiquitination involves at least three classes of enzymes: ubiquitin-activating enzymes, ubiquitin-conjugating enzymes, and ubiquitin-protein ligases. This gene encodes a member of the E2 ubiquitin-conjugating enzyme family. The encoded protein is required for the destruction of mitotic cyclins and for cell cycle progression, and may be involved in cancer progression. Multiple transcript variants encoding different isoforms have been found for this gene. Pseudogenes of this gene have been defined on chromosomes 4, 14, 15, 18, and 19. [provided by RefSeq, Aug 2013]

Locus ID:

11065

MW:

8.5