

## Product datasheet for SC202703

### TNNT3 (NM\_006757) Human 3' UTR Clone

#### Product data:

Product Type:	3' UTR Clones
Product Name:	TNNT3 (NM_006757) Human 3' UTR Clone
Symbol:	TNNT3
Synonyms:	beta-TnTF; DA2B2; TNTF
Mammalian Cell Selection:	Neomycin
Vector:	pMirTarget (PS100062)
ACCN:	NM_006757
Insert Size:	240 bp
Insert Sequence:	<p>&gt;SC202703 3'UTR clone of NM_006757</p> <p>The sequence shown below is from the reference sequence of NM_006757. The complete sequence of this clone may contain minor differences, such as SNPs.</p> <p>Blue=Stop Codon Red=Cloning site</p>

```

GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAGGCCAAGAAGGGCGGAAAGATCGCCGTG
TAACAATTGGCAGAGCTCAGAATTCAAACGATCGCC
AAGGGCAAAGTCGGCGGCGCTGGAAGTAGAGAGGCCAGAAAGGCCCTCGAGGCAGAGACCCTCCGCC
CTCTTGACACACAGGGCCGCTCGTGGGACTCCACATCCTCCAGCCCCACAATCCTGTCAGGGGCTCCC
TGACAGTCCTGGGGTGGAGAGGCCATCCCGGGCGTCCCCCGCTCTGTGTCCTTGCTGCCTTCATCC
CCTGGGGCCTGTGAATAAAGCTGCAGAACCCCC
ACGCGTAAGCGGCCGCGCATCTAGATTGAAGAAAATGACCGACCAAGCGACGCCAACCTGCCATCA
CGAGATTCGATTCCACCGCCGCTTCTATGAAAGG
  
```

Restriction Sites:	SgfI-MluI
OTI Disclaimer:	Our molecular clone sequence data has been matched to the sequence identifier above as a point of reference. Note that the complete sequence of this clone is largely the same as the reference sequence but may contain minor differences, e.g., single nucleotide polymorphisms (SNPs).
Components:	The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The package also includes 100 pmols of both the corresponding 5' and 3' vector primers in separate vials.
RefSeq:	<a href="#">NM_006757.4</a>


[View online »](#)

**Summary:**

The binding of Ca(2+) to the trimeric troponin complex initiates the process of muscle contraction. Increased Ca(2+) concentrations produce a conformational change in the troponin complex that is transmitted to tropomyosin dimers situated along actin filaments. The altered conformation permits increased interaction between a myosin head and an actin filament which, ultimately, produces a muscle contraction. The troponin complex has protein subunits C, I, and T. Subunit C binds Ca(2+) and subunit I binds to actin and inhibits actin-myosin interaction. Subunit T binds the troponin complex to the tropomyosin complex and is also required for Ca(2+)-mediated activation of actomyosin ATPase activity. There are 3 different troponin T genes that encode tissue-specific isoforms of subunit T for fast skeletal-, slow skeletal-, and cardiac-muscle. This gene encodes fast skeletal troponin T protein; also known as troponin T type 3. Alternative splicing results in multiple transcript variants encoding additional distinct troponin T type 3 isoforms. A developmentally regulated switch between fetal/neonatal and adult troponin T type 3 isoforms occurs. Additional splice variants have been described but their biological validity has not been established. Mutations in this gene may cause distal arthrogryposis multiplex congenita type 2B (DA2B). [provided by RefSeq, Oct 2009]

**Locus ID:**

7140

**MW:**

8.3