

## Product datasheet for **SC202755**

### PSMD2 (NM\_002808) Human 3' UTR Clone

#### Product data:

Product Type:	3' UTR Clones
Product Name:	PSMD2 (NM_002808) Human 3' UTR Clone
Vector:	pMirTarget (PS100062)
Symbol:	PSMD2
Synonyms:	P97; RPN1; S2; TRAP2
ACCN:	NM_002808
Insert Size:	190 bp
Insert Sequence:	>SC202755 3'UTR clone of NM_002808 The sequence shown below is from the reference sequence of NM_002808. The complete sequence of this clone may contain minor differences, such as SNPs. <b>Blue</b> =Stop Codon <b>Red</b> =Cloning site  GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAAGCCAAGAAGGGCGGAAAGATCGCCGTG TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC CTTCGGAAGAACCCCAATTATGATCTCTAAGTGACCACCAGGGGCTCTGAACTGCAGCTGATGTTATCA GCAGGCCATGCATCCTGCTGCCAAGGGTGGACACGGCTGCAGACTCTGGGGGAATTGTCGCCTCCTGC TCTTTTGTACTGAGTGAGATAAGGTTGTTCAATAAAGACTTTTATCCCCAA <b>ACGCGT</b> AAGCGGCCGCGCATCTAGATTGAAAGAAAATGACCGACCAAGCGACGCCCAACCTGCCATCA CGAGATTCGATTCCACCGCCCTTCTATGAAAGG
Restriction Sites:	Sgfl-Mlul
OTI Disclaimer:	Our molecular clone sequence data has been matched to the sequence identifier above as a point of reference. Note that the complete sequence of this clone is largely the same as the reference sequence but may contain minor differences , e.g., single nucleotide polymorphisms (SNPs).
Components:	The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The package also includes 100 pmols of both the corresponding 5' and 3' vector primers in separate vials.
RefSeq:	<a href="#">NM_002808.5</a>



[View online »](#)

**Summary:**

The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes one of the non-ATPase subunits of the 19S regulator lid. In addition to participation in proteasome function, this subunit may also participate in the TNF signalling pathway since it interacts with the tumor necrosis factor type 1 receptor. A pseudogene has been identified on chromosome 1. Alternative splicing results in multiple transcript variants of this gene. [provided by RefSeq, Jul 2013]

**Locus ID:**

5708

**MW:**

6.9