

Product datasheet for **SC203453**

HNMT (NM_001024075) Human 3' UTR Clone

Product data:

Product Type:	3' UTR Clones
Product Name:	HNMT (NM_001024075) Human 3' UTR Clone
Vector:	pMirTarget (PS100062)
Symbol:	HNMT
Synonyms:	HMT; HNMT-S1; HNMT-S2; MRT51
ACCN:	NM_001024075
Insert Size:	248 bp
Insert Sequence:	>SC203453 3'UTR clone of NM_001024075 The sequence shown below is from the reference sequence of NM_001024075. The complete sequence of this clone may contain minor differences, such as SNPs. Blue=Stop Codon Red=Cloning site GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAAGCCAAGAAGGGCGGAAAGATCGCCGTG TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC TCATTCCTAGTTTCCTTCATCCTCTTTTAAAGTGCATCTTTTCCATGAAGCCTACTAGATCACACAGTT GAAAATTGCAAAGTCTTACACCCGAGCTTCCCCTTTCTTTGTTGCTTTGGTTTTCTCCGTGGGACTTC CCACCATCCGTAATAATTAGGTGTTTTTGCCTGTTTTATTCACTGCTACGTCTCCAATGGTTAGAACCTG ACATTTAGTGGTGCTCAATAATATTTGTTGAAGTAATGTA ACGCGTAAGCGGCCGCGCATCTAGATTGAAAGAAAATGACCGACCAAGCGACGCCAACCTGCCATCA CGAGATTCGATTCCACCGCCGCTTCTATGAAAGG
Restriction Sites:	SgfI-MluI
OTI Disclaimer:	Our molecular clone sequence data has been matched to the sequence identifier above as a point of reference. Note that the complete sequence of this clone is largely the same as the reference sequence but may contain minor differences, e.g., single nucleotide polymorphisms (SNPs).
Components:	The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The package also includes 100 pmols of both the corresponding 5' and 3' vector primers in separate vials.
RefSeq:	NM_001024075.3



[View online »](#)

Summary:

In mammals, histamine is metabolized by two major pathways: N(tau)-methylation via histamine N-methyltransferase and oxidative deamination via diamine oxidase. This gene encodes the first enzyme which is found in the cytosol and uses S-adenosyl-L-methionine as the methyl donor. In the mammalian brain, the neurotransmitter activity of histamine is controlled by N(tau)-methylation as diamine oxidase is not found in the central nervous system. A common genetic polymorphism affects the activity levels of this gene product in red blood cells. Multiple alternatively spliced transcript variants that encode different proteins have been found for this gene. [provided by RefSeq, Jul 2008]

Locus ID:

3176

MW:

9.2