

## **Product datasheet for SC203655**

## **SGCE (NM 001099401) Human 3' UTR Clone**

**Product data:** 

**Product Type:** 3' UTR Clones

**Product Name:** SGCE (NM\_001099401) Human 3' UTR Clone

**Vector:** pMirTarget (PS100062)

Symbol: SGCE

Synonyms: DYT11; epsilon-SG; ESG

**ACCN:** NM\_001099401

**Insert Size:** 549 bp

The sequence shown below is from the reference sequence of NM\_001099401. The complete

sequence of this clone may contain minor differences, such as SNPs.

Blue=Stop Codon Red=Cloning site

GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAGGCCAAGAAGGGCGGAAAGATCGCCGTG

TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC

Restriction Sites: Sgfl-Mlul

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the sequence identifier above as a

point of reference. Note that the complete sequence of this clone is largely the same as the

reference sequence but may contain minor differences, e.g., single nucleotide

polymorphisms (SNPs).

**Components:** The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The

package also includes 100 pmols of both the corresponding 5' and 3' vector primers in

separate vials.

**RefSeg:** NM 001099401.2



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## SGCE (NM\_001099401) Human 3' UTR Clone - SC203655

Summary: This gene encodes the epsilon member of the sarcoglycan family. Sarcoglycans are

transmembrane proteins that are components of the dystrophin-glycoprotein complex, which link the actin cytoskeleton to the extracellular matrix. Unlike other family members which are predominantly expressed in striated muscle, the epsilon sarcoglycan is more broadly expressed. Mutations in this gene are associated with myoclonus-dystonia syndrome. This gene is imprinted, with preferential expression from the paternal allele. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. A pseudogene associated with this gene is located on chromosome 2. [provided by RefSeq, Oct 2016]

**Locus ID:** 8910

MW: 21.4