

## Product datasheet for **SC203801**

### SPHK1 (NM\_001142602) Human 3' UTR Clone

#### Product data:

Product Type:	3' UTR Clones
Product Name:	SPHK1 (NM_001142602) Human 3' UTR Clone
Vector:	pMirTarget (PS100062)
Symbol:	SPHK1
Synonyms:	SPHK
ACCN:	NM_001142602
Insert Size:	299 bp
Insert Sequence:	>SC203801 3'UTR clone of NM_001142602 The sequence shown below is from the reference sequence of NM_001142602. The complete sequence of this clone may contain minor differences, such as SNPs. <b>Blue</b> =Stop Codon <b>Red</b> =Cloning site  GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAAGCCAAGAAGGGCGGAAAGATCGCCGTG TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC CAGATGCCACCGCCAGAAGAGCCCTTATGACCCCTGGGCCGCGCTGTGCCTTAGTGTCTACTTGCAGGA CCCTTCTCCTTCCCTAGGGCTGCAGGGCCTGTCCACAGCTCCTGTGGGGTGGAGGAGACTCCTCTGG AGAAGGGTGAGAAGGTGGAGGCTATGCTTTGGGGGACAGGCCAGAATGAAGTCCGGGTGAGGAGCCC AGCTGGCTGGGCCAGCTGCCTATGTAAGGCCTTCTAGTTTGTCTGAGACCCCCACCCACGAACCAA ATCCAAATAAAGTGACATTCCCA <b>ACGCGT</b> AAGCGGCCGCGGCATCTAGATTGAAGAAAATGACCGACCAAGCGACGCCAACCTGCCATCA CGAGATTCGATTCCACCGCCCTTCTATGAAAGG
Restriction Sites:	Sgfl-MluI
OTI Disclaimer:	Our molecular clone sequence data has been matched to the sequence identifier above as a point of reference. Note that the complete sequence of this clone is largely the same as the reference sequence but may contain minor differences, e.g., single nucleotide polymorphisms (SNPs).
Components:	The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The package also includes 100 pmols of both the corresponding 5' and 3' vector primers in separate vials.
RefSeq:	<a href="#">NM_001142602.2</a>



[View online »](#)

**Summary:**

The protein encoded by this gene catalyzes the phosphorylation of sphingosine to form sphingosine-1-phosphate (S1P), a lipid mediator with both intra- and extracellular functions. Intracellularly, S1P regulates proliferation and survival, and extracellularly, it is a ligand for cell surface G protein-coupled receptors. This protein, and its product S1P, play a key role in TNF-alpha signaling and the NF-kappa-B activation pathway important in inflammatory, antiapoptotic, and immune processes. Phosphorylation of this protein alters its catalytic activity and promotes its translocation to the plasma membrane. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Sep 2017]

**Locus ID:** 8877

**MW:** 11.3