

Product datasheet for SC203804

SPHK1 (NM_182965) Human 3' UTR Clone

Product data:

Product Type: 3' UTR Clones
Product Name: SPHK1 (NM_182965) Human 3' UTR Clone
Symbol: SPHK1
Synonyms: SPHK
Mammalian Cell Selection: Neomycin
Vector: pMirTarget (PS100062)
ACCN: NM_182965
Insert Size: 299 bp
Insert Sequence: >SC203804 3'UTR clone of NM_182965
 The sequence shown below is from the reference sequence of NM_182965. The complete sequence of this clone may contain minor differences, such as SNPs.
 Blue=Stop Codon Red=Cloning site

```

GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAGGCCAAGAAGGGCGGAAAGATCGCCGTG
TAACAATTGGCAGAGCTCAGAATTCAAACGATCGCC
CAGATGCCACCGCCAGAAGAGCCCTTAGACCCCTGGGCCGCGCTGTGCCTTAGTGTCTACTTGCAGGA
CCCTTCCTCCTCCCTAGGGCTGCAGGGCCTGTCCACAGCTCCTGTGGGGGTGGAGGAGACTCCTCTGG
AGAAGGGTGAGAAGGTGGAGGCTATGCTTTGGGGGGACAGGCCAGAATGAAGTCTGGGTCAGGAGCCC
AGCTGGCTGGGCCAGCTGCCTATGTAAGGCCTTCTAGTTTGTCTGAGACCCCCACCCACGAACCAA
ATCCAAATAAAGTGACATTCCCA
ACGCGTAAGCGGCCGCGCATCTAGATTGGAAGAAATGACCGACCAAGCGACGCCAACCTGCCATCA
CGAGATTCGATTCCACCGCCGCTTCTATGAAAGG
  
```

Restriction Sites: SgfI-MluI
OTI Disclaimer: Our molecular clone sequence data has been matched to the sequence identifier above as a point of reference. Note that the complete sequence of this clone is largely the same as the reference sequence but may contain minor differences, e.g., single nucleotide polymorphisms (SNPs).
Components: The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The package also includes 100 pmols of both the corresponding 5' and 3' vector primers in separate vials.
RefSeq: [NM_182965.3](#)


[View online »](#)

Summary:

The protein encoded by this gene catalyzes the phosphorylation of sphingosine to form sphingosine-1-phosphate (S1P), a lipid mediator with both intra- and extracellular functions. Intracellularly, S1P regulates proliferation and survival, and extracellularly, it is a ligand for cell surface G protein-coupled receptors. This protein, and its product S1P, play a key role in TNF- α signaling and the NF- κ B activation pathway important in inflammatory, antiapoptotic, and immune processes. Phosphorylation of this protein alters its catalytic activity and promotes its translocation to the plasma membrane. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Sep 2017]

Locus ID:

8877

MW:

11.3