

Product datasheet for SC204175

1 Todact datasticct for Sc204175

RPL41 (NM_001035267) Human 3' UTR Clone

Product data:

Product Type: 3' UTR Clones

Product Name: RPL41 (NM 001035267) Human 3' UTR Clone

Symbol: RPL41
Synonyms: L41

Mammalian Cell

Selection:

Neomycin

Vector: pMirTarget (PS100062)

ACCN: NM_001035267

Insert Size: 473 bp

Insert Sequence: >SC204175 3'UTR clone of NM_001035267

The sequence shown below is from the reference sequence of NM_001035267. The complete

sequence of this clone may contain minor differences, such as SNPs.

Blue=Stop Codon Red=Cloning site

GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAGGCCAAGAAGGGCGGAAAGATCGCCGTG

TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC

GCTATGGAGGGTATCAAAGCTATAAAGGCAACAGCCCGGGTTACGTGGTGTGGTTTGCA

CGAGATTTCGATTCCACCGCCGCCTTCTATGAAAGG

Restriction Sites: Sgfl-Mlul

OTI Disclaimer: Our molecular clone sequence data has been matched to the sequence identifier above as a

point of reference. Note that the complete sequence of this clone is largely the same as the

reference sequence but may contain minor differences, e.g., single nucleotide

polymorphisms (SNPs).

Components: The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The

package also includes 100 pmols of both the corresponding 5' and 3' vector primers in

separate vials.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



RefSeq: <u>NM 001035267.2</u>

Summary: Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and

a large 60S subunit. Together these subunits are composed of 4 RNA species and

approximately 80 structurally distinct proteins. This gene encodes a ribosomal protein that is a component of the 60S subunit. The protein, which shares sequence similarity with the yeast ribosomal protein YL41, belongs to the L41E family of ribosomal proteins. It is located in the cytoplasm. The protein can interact with the beta subunit of protein kinase CKII and can stimulate the phosphorylation of DNA topoisomerase II-alpha by CKII. Two alternative splice variants have been identified, both encoding the same protein. As is typical for genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene

dispersed through the genome. [provided by RefSeq, Jul 2008]

Locus ID: 6171 **MW:** 17.6