

Product datasheet for SC204248

HLA-DRB4 (NM 021983) Human 3' UTR Clone

Product data:

Product Type: 3' UTR Clones

Product Name: HLA-DRB4 (NM_021983) Human 3' UTR Clone

Vector: pMirTarget (PS100062)

Symbol: HLA-DRB4

Synonyms: DR4; DRB4; HLA-DR4B; HLA-DRB4*

ACCN: NM_021983

Insert Size: 351 bp

Insert Sequence: >SC204248 3'UTR clone of NM_021983

The sequence shown below is from the reference sequence of NM_021983. The complete

sequence of this clone may contain minor differences, such as SNPs.

Blue=Stop Codon Red=Cloning site

GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAGGCCAAGAAGGGCGGAAAGATCGCCGTG

TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC

GGACTTCAGCCAACAGGACTCTTGAGCTGAAGTGCAGATGACCACATTCAAGGAAGAACCTTCTGCCCC
AGCTTTGCAAGATGAAAAGCTTTCCCACTTGGCTCTTATTCTTCCACAAGAGCTTTGTCAGGACCAGGT
TGTTACTGGTTCAGCAACTCTGCAGAAAATGTCCTCCCTTGTGGCTTCCTTAGCTCCTGTTCTTGGCCT
GAAGCCTCACAGCTTTGATGGCAGTGCCTCATCTTCAACTTTTTGTGCTTCCCTTTACCTAAACTGTCCT
GCCTCCCGTGCATCTGTACTCCCCTTGTGCCACACATTGCATTATTAAATGTTTCTCAAACATGGAGTT

AAAAA

CGAGATTTCGATTCCACCGCCGCCTTCTATGAAAGG

Restriction Sites: Sgfl-Mlul

OTI Disclaimer: Our molecular clone sequence data has been matched to the sequence identifier above as a

point of reference. Note that the complete sequence of this clone is largely the same as the

reference sequence but may contain minor differences, e.g., single nucleotide

polymorphisms (SNPs).

Components: The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The

package also includes 100 pmols of both the corresponding 5' and 3' vector primers in

separate vials.

RefSeg: NM 021983.4



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



ORIGENE

Summary: HLA-DRB4 belongs to the HLA class II beta chain paralogues. This class II molecule is a

heterodimer consisting of an alpha (DRA) and a beta (DRB) chain, both anchored in the membrane. It plays a central role in the immune system by presenting peptides derived from extracellular proteins. Class II molecules are expressed in antigen presenting cells. The beta chain is approximately 26-28 kDa and its gene contains 6 exons. Exon one encodes the leader

peptide, exons 2 and 3 encode the two extracellular domains, exon 4 encodes the

transmembrane domain and exon 5 encodes the cytoplasmic tail. Within the DR molecule the beta chain contains all the polymorphisms specifying the peptide binding specificities. Typing for these polymorphisms is routinely done for bone marrow and kidney transplantation.

There are multiple pseudogenes of this gene. [provided by RefSeq, Feb 2020]

Locus ID: 3126

MW: 12.9