

Product datasheet for SC204691

PABPC4 (NM_003819) Human 3' UTR Clone

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	3' UTR Clones
Product Name:	PABPC4 (NM_003819) Human 3' UTR Clone
Symbol:	PABPC4
Synonyms:	APP-1; APP1; iPABP; PABP4
Mammalian Cell Selection:	Neomycin
Vector:	pMirTarget (PS100062)
ACCN:	NM_003819
Insert Size:	350 bp
Insert Sequence:	<pre>>SC204691 3'UTR clone of NM_003819 The sequence shown below is from the reference sequence of NM_003819. The complete sequence of this clone may contain minor differences, such as SNPs. Blue=Stop Codon Red=Cloning site GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAGGCCAAGAAGGGCGGAAAGATCGCCGTG TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC GTGGGCGCTGTTGCTGCTGCTACCTCTTAGACAAGGAAAAACCGATTCAAAAGCCAAATAACCCCTTAT GGAATTCAACTCAAGGTTTGAAGACTTCCTAGCTTGTCCTATGGACCTCAACACCAAGGATTACAAATT GCAAATTTAATAGGTCATTTGTATCAAAAGGTCAATTATGAAGCCCAAGAATTTTTCAATTATACGA ATATGTTCTTTGGGTTCTGCTGTGGCCCAGACAGTGTTAACTTTTTTTT</pre>
Restriction Sites:	Sgfl-Mlul
OTI Disclaimer:	Our molecular clone sequence data has been matched to the sequence identifier above as a point of reference. Note that the complete sequence of this clone is largely the same as the reference sequence but may contain minor differences , e.g., single nucleotide polymorphisms (SNPs).
Components:	The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The package also includes 100 pmols of both the corresponding 5' and 3' vector primers in separate vials.



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	PABPC4 (NM_003819) Human 3' UTR Clone – SC204691
RefSeq:	<u>NM 003819.4</u>
Summary:	Poly(A)-binding proteins (PABPs) bind to the poly(A) tail present at the 3-prime ends of most eukaryotic mRNAs. PABPC4 or IPABP (inducible PABP) was isolated as an activation-induced T- cell mRNA encoding a protein. Activation of T cells increased PABPC4 mRNA levels in T cells approximately 5-fold. PABPC4 contains 4 RNA-binding domains and proline-rich C terminus. PABPC4 is localized primarily to the cytoplasm. It is suggested that PABPC4 might be necessary for regulation of stability of labile mRNA species in activated T cells. PABPC4 was also identified as an antigen, APP1 (activated-platelet protein-1), expressed on thrombin- activated rabbit platelets. PABPC4 may also be involved in the regulation of protein translation in platelets and megakaryocytes or may participate in the binding or stabilization of polyadenylates in platelet dense granules. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Oct 2008]
Locus ID:	8761
MW:	12.9

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US