

## Product datasheet for SC205332

### BAD (NM\_032989) Human 3' UTR Clone

#### Product data:

Product Type:	3' UTR Clones
Product Name:	BAD (NM_032989) Human 3' UTR Clone
Symbol:	BAD
Synonyms:	BBC2; BCL2L8
Mammalian Cell Selection:	Neomycin
Vector:	pMirTarget (PS100062)
ACCN:	NM_032989
Insert Size:	409 bp
Insert Sequence:	>SC205332 3'UTR clone of NM_032989 The sequence shown below is from the reference sequence of NM_032989. The complete sequence of this clone may contain minor differences, such as SNPs. <b>Blue</b> =Stop Codon <b>Red</b> =Cloning site
Restriction Sites:	Sgfl-MluI
OTI Disclaimer:	Our molecular clone sequence data has been matched to the sequence identifier above as a point of reference. Note that the complete sequence of this clone is largely the same as the reference sequence but may contain minor differences , e.g., single nucleotide polymorphisms (SNPs).
Components:	The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The package also includes 100 pmols of both the corresponding 5' and 3' vector primers in separate vials.

```
GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAAGCCAAGAAGGGCGGAAAGATCGCCGTG
TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC
GGCAGGGGAAGCTCCGCCCTCCAGTGACCTTCGCTCCACATCCCGAACTCCACCCGTTCCCACTG
CCCTGGGCAGCCATCTTGAATATGGGCGGAAGTACTTCCCTCAGGCCTATGCAAAAAGAGGATCCGTGC
TGTCTCCTTTGGAGGGAGGGCTGACCCAGATTCCCTTCCGGTGCCTGTGAAGCCACGGAAGGCTTGGTC
CCATCGGAAGTTTTGGGTTTTCCGCCACAGCCCGGAAGTGGCTCCGTGGCCCGCCCTCAGGCTCC
GGGCTTTCCCCAGGCGCTGCGCTAAGTCGCGAGCCAGGTTAAACCGTTGCGTCAACGGGACCCGAGC
CCCCGCGATGCCCTGGGGCCGTGCTCACTACCAAATGTTAATAAAGCCCGCTGTGCGCCG
ACGCGTAAGCGGCCGCGCATCTAGATTGAAGAAAATGACCGACCAAGCGACGCCCAACCTGCCATCA
CGAGATTCGATTCCACCGCCGCTTCTATGAAAGG
```



[View online »](#)

RefSeq: [NM\\_032989.3](#)

**Summary:** The protein encoded by this gene is a member of the BCL-2 family. BCL-2 family members are known to be regulators of programmed cell death. This protein positively regulates cell apoptosis by forming heterodimers with BCL-xL (B-cell lymphoma-extra large) and BCL-2, and reversing their death repressor activity. Proapoptotic activity of this protein is regulated through its phosphorylation. Protein kinases AKT and MAP kinase, as well as protein phosphatase calcineurin were found to be involved in the regulation of this protein. Alternative splicing of this gene results in two transcript variants which encode the same isoform. [provided by RefSeq, Dec 2019]

**Locus ID:** 572

**MW:** 14.4