

Product datasheet for SC205450

IL5RA (NM_175727) Human 3' UTR Clone

Product data:

Product Type:	3' UTR Clones
Product Name:	IL5RA (NM_175727) Human 3' UTR Clone
Symbol:	IL5RA
Synonyms:	CD125; CDw125; HSIL5R3; IL5R
Mammalian Cell Selection:	Neomycin
Vector:	pMirTarget (PS100062)
ACCN:	NM_175727
Insert Size:	420 bp
Insert Sequence:	<p>>SC205450 3' UTR clone of NM_175727</p> <p>The sequence shown below is from the reference sequence of NM_175727. The complete sequence of this clone may contain minor differences, such as SNPs. Red=Cloning site Blue=Stop Codon</p>

CAATTGGCAGAGCTCAGAATTCAA**GCGATCGC**

AGTGAGTGGAGCCAACCTATTTATGTGGGTAAG**TAG**CTTATGTTTATTTTACATTGGCAGCCTTCCTTGT
 GATCAAAAAGGTAATCCAGAAACGTACCGTTCACTCGTGGGTCTTAAATGGTTTCATATCTCTATT
 GTGACTAATTTTCTCTCGGTCTACTGCCTTTTCAATCAGGAATAGATTTGCCATGAAGCCAGTGAAGTTT
 TTAAGTGTCTAGGCTTCTCATTAGCGCCAACCTCCTAGACCTGGTGCCTGTTTTTTTCCAAGTTTGT
 TTCTACTTCTATCCATTTTTTAAATTAACCTTTTATTTTGAATAATTATCACACTCACAAGCTGTGGG
 AAGAAATAATAGAGATCCTGTGTCTTTTCATCCAGTTTCTCAAGGGTAACATCTTACAAAATATAG

ACGCGTAAGCGGCCGCGGCATCTAGATTGAAGAAAATGACCG

Restriction Sites:	Sgfi-MluI
OTI Disclaimer:	Our molecular clone sequence data has been matched to the sequence identifier above as a point of reference. Note that the complete sequence of this clone is largely the same as the reference sequence but may contain minor differences , e.g., single nucleotide polymorphisms (SNPs).
Components:	The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The package also includes 100 pmols of both the corresponding 5' and 3' vector primers in separate vials.


[View online »](#)

RefSeq: NM_175727.2

Summary: The protein encoded by this gene is an interleukin 5 specific subunit of a heterodimeric cytokine receptor. The receptor is comprised of a ligand specific alpha subunit and a signal transducing beta subunit shared by the receptors for interleukin 3 (IL3), colony stimulating factor 2 (CSF2/GM-CSF), and interleukin 5 (IL5). The binding of this protein to IL5 depends on the beta subunit. The beta subunit is activated by the ligand binding, and is required for the biological activities of IL5. This protein has been found to interact with syndecan binding protein (syntenin), which is required for IL5 mediated activation of the transcription factor SOX4. Several alternatively spliced transcript variants encoding four distinct isoforms have been reported. [provided by RefSeq, Jul 2011]

Locus ID: 3568