

Product datasheet for **SC205487**

SKAP55 (SKAP1) (NM_003726) Human 3' UTR Clone

Product data:

Product Type:	3' UTR Clones
Symbol:	SKAP55
Synonyms:	HEL-S-8lp; SCAP1; SKAP55
Mammalian Cell	Neomycin
Selection:	
Vector:	pMirTarget (PSI00062)
ACCN:	NM_003726
Insert Size:	412 bp
Insert Sequence:	<p>>SC205487 3'UTR clone of NM_003726</p> <p>The sequence shown below is from the reference sequence of NM_003726. The complete sequence of this clone may contain minor differences, such as SNPs.</p> <p>Blue=Stop Codon Red=Cloning site</p> <pre> GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAAGCCAAGAAGGGCGGAAAGATCGCCGTG TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC ACCACTGCCTTTGAAGTGAAGAAAGATGAACCCAGGAAATATATTCTTCCCTCTCTCCTGCCTTTAT GAGGAAACTGATCATCAAAAGTTCCCACTCCCTACTTCTGCCACCCACCAACGCCCTTGGACTCCTCTC TTTGCTGAAGAGACCCAAGTCTCTTGACACCTCAGAGTGACTGTAAGCTACCAGTAAGACAAGTGGGAA GAGGCACGTTTCATCAAACTGTGTTACTAAACAGCCTAGTCATAGCTCATCCCCATCTCTAAATGTGTCC ACACAACCACATCTGCCTTTTCCACAAGCTTTTCAAAAGAAGGTGAGAGAGAAGGAAACCTTGGGAGG AGGACATTACTGGTTGTTCTGGCTGGTTTGAAGACACAAATAAACTTGGGATGTGGTTCTTGCCA ACGCGTAAGCGGCCGCGGCATCTAGATTCAAGAAAAATGACCGACCAAGCGACGCCCAACCTGCCATCA CGAGATTTTCGATTCCACCGCCGCTTCTATGAAAGG </pre>
Restriction Sites:	SgfI-MluI
OTI Disclaimer:	Our molecular clone sequence data has been matched to the sequence identifier above as a point of reference. Note that the complete sequence of this clone is largely the same as the reference sequence but may contain minor differences, e.g., single nucleotide polymorphisms (SNPs).



Components:	The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The package also includes 100 pmols of both the corresponding 5' and 3' vector primers in separate vials.
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	<u>NM_003726.4</u>
Summary:	This gene encodes a T cell adaptor protein, a class of intracellular molecules with modular domains capable of recruiting additional proteins but that exhibit no intrinsic enzymatic activity. The encoded protein contains a unique N-terminal region followed by a PH domain and C-terminal SH3 domain. Along with the adhesion and degranulation-promoting adaptor protein, the encoded protein plays a critical role in inside-out signaling by coupling T-cell antigen receptor stimulation to the activation of integrins. [provided by RefSeq, Jul 2008]
Locus ID:	8631
MW:	15.5