

Product datasheet for **SC206365**

UBASH3A (NM_018961) Human 3' UTR Clone

Product data:

Product Type:	3' UTR Clones
Product Name:	UBASH3A (NM_018961) Human 3' UTR Clone
Symbol:	UBASH3A
Synonyms:	CLIP4; STS-2; TULA; TULA-1
Mammalian Cell Selection:	Neomycin
Vector:	pMirTarget (PS100062)
ACCN:	NM_018961
Insert Size:	520 bp
Insert Sequence:	>SC206365 3'UTR clone of NM_018961 The sequence shown below is from the reference sequence of NM_018961. The complete sequence of this clone may contain minor differences, such as SNPs. Blue =Stop Codon Red =Cloning site

```
GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAAGCCAAGAAGGGCGGAAAGATCGCCGTG
TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC
AACTGGAGGAACTGGATCTCAGGCAACTTGAGAGCCACGGTGATGTTGTCATAACCTCAGAGTGGAGAGG
CAGAAACCATGTGCAGAGGCTGGGAGATGCTGCTGTTCCAGAGGCGTCTTAGTCTACCCAATGTGAT
TTGTAGAAGCACGAGACGCACTTTTATATCCCGGAATATTTCCCTCCGGCTTTCGCCTTTGTAACCCC
ATCTGTGGACCCATCGTCCACCAGCCAGCTGCGGGGAGCACAGGGCAGGTGGCTGGGTGAGGATGCCG
CCCTGCAGCATGTACACCGAGTGTCTGCAGCTGGGGACACAACCTGCCCGGACTCTAACTCCAGGAAT
TAAAGACTCACACACGAAGGATCTAACCATTCTATTTCCATGGTCTAATCATTAAATTCCCAATC
GTTTTTCTTTTTCTGGGTCCATCACTCTTTAGCCATATCCACATGGGCTAAAACAGGTGTAATAGTCA
ATAAAATGGTCCCAGAAAACCATCGACCAGGTCAAAA
ACGCGTAAGCGGCCGCGGCATCTAGATTCAAGAAAAATGACCGACCAAGCGACGCCCAACCTGCCATCA
CGAGATTCGATTCCACCGCCGCTTCTATGAAAGG
```

Restriction Sites:	Sgfl-MluI
OTI Disclaimer:	Our molecular clone sequence data has been matched to the sequence identifier above as a point of reference. Note that the complete sequence of this clone is largely the same as the reference sequence but may contain minor differences , e.g., single nucleotide polymorphisms (SNPs).



[View online »](#)

Components:	The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The package also includes 100 pmols of both the corresponding 5' and 3' vector primers in separate vials.
RefSeq:	NM_018961.4
Summary:	This gene encodes one of two family members belonging to the T-cell ubiquitin ligand (TULA) family. Both family members can negatively regulate T-cell signaling. This family member can facilitate growth factor withdrawal-induced apoptosis in T cells, which may occur via its interaction with AIF, an apoptosis-inducing factor. Alternative splicing of this gene results in multiple transcript variants. [provided by RefSeq, Aug 2011]
Locus ID:	53347
MW:	19.8