

Product datasheet for SC206417

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Parathyroid hormone related protein (PTHLH) (NM 198965) Human 3' UTR Clone

Product data:

Product Type: 3' UTR Clones

Product Name: Parathyroid hormone related protein (PTHLH) (NM 198965) Human 3' UTR Clone

Vector: pMirTarget (PS100062)

Symbol: PTHLH

Synonyms: BDE2; HHM; PLP; PTHR; PTHRP

ACCN: NM_198965

Insert Size: 505 bp

Insert Sequence: >SC206417 3'UTR clone of NM_198965

The sequence shown below is from the reference sequence of NM_198965. The complete

sequence of this clone may contain minor differences, such as SNPs.

Blue=Stop Codon Red=Cloning site

GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAGGCCAAGAAGGGCGGAAAGATCGCCGTG

TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC

ATATAATCTAATTACATCATGA

CGAGATTTCGATTCCACCGCCGCCTTCTATGAAAGG

Restriction Sites: Sgfl-Mlul

OTI Disclaimer: Our molecular clone sequence data has been matched to the sequence identifier above as a

point of reference. Note that the complete sequence of this clone is largely the same as the

reference sequence but may contain minor differences, e.g., single nucleotide

polymorphisms (SNPs).

Components: The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The

package also includes 100 pmols of both the corresponding 5' and 3' vector primers in

separate vials.

RefSeg: NM 198965.2





Parathyroid hormone related protein (PTHLH) (NM_198965) Human 3' UTR Clone - SC206417

Summary: The protein encoded by this gene is a member of the parathyroid hormone family. This

hormone, via its receptor, PTHR1, regulates endochondral bone development and epithelial-mesenchymal interactions during the formation of the mammary glands and teeth. It is responsible for most cases of humoral hypercalcemia of malignancy, and mutations in this gene are associated with brachydactyly type E2 (BDE2). Alternatively spliced transcript variants have been found for this gene. There is also evidence for alternative translation initiation from non-AUG (CUG and GUG) start sites, downstream of the initiator AUG codon, resulting in nuclear forms of this hormone. [provided by RefSeq, Nov 2013]

Locus ID: 5744

MW: 20.1