

Product datasheet for SC206666

CXCR3 (NM_001504) Human 3' UTR Clone

Product data:

| | |
|---------------------------|--|
| Product Type: | 3' UTR Clones |
| Product Name: | CXCR3 (NM_001504) Human 3' UTR Clone |
| Symbol: | CXCR3 |
| Synonyms: | CD182; CD183; CKR-L2; CMKAR3; GPR9; IP10-R; Mig-R; MigR |
| Mammalian Cell Selection: | Neomycin |
| Vector: | pMirTarget (PS100062) |
| ACCN: | NM_001504 |
| Insert Size: | 476 bp |
| Insert Sequence: | >SC206666 3'UTR clone of NM_001504 The sequence shown below is from the reference sequence of NM_001504. The complete sequence of this clone may contain minor differences, such as SNPs. Blue =Stop Codon Red =Cloning site |

```
GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAAGCCAAGAAGGGCGGAAAGATCGCCGTG
TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC
ACCTCAGAGGCCCTCTACTCGGGCTTGTGAGGCCGGAATCCGGGCTCCCTTTTGGCCACAGTCTGACT
TCCCCGATTCCAGGCTCCTCCCTCCCTCTGCCGGCTCTGGCTCTCCCAATATCCTCGCTCCCGGGAC
TACTGGCAGCCCAGCACCACCAGGTCTCCCGGAAGCCACCCTCCAGCTCTGAGGACTGCACCATT
GCTGCTCCTTAGCTGCCAAGCCCATCCTGCCGCCGAGGTGGCTGCCTGGAGCCCCACTGCCCTTCTC
ATTGGAAACTAAAATTTCATCTTCCCAAGTGCGGGGAGTACAAGGCATGGCGTAGAGGGTGTGCC
CATGAAGCCACAGCCCAGGCCTCAGCTCAGCAGTGACTGTGGCCATGGTCCCAAGACCTTATATTT
GCTTTTTATTTTATGTCTAAAATCCTGCTTAAAACCTTTTCAATAAAACAAGATCGTCAGGA
ACGCGTAAGCGGCCGCGCATCTAGATTGAAGAAAATGACCGACCAAGCGACGCCAACCTGCCATCA
CGAGATTCGATTCCACCGCCGCTTCTATGAAAGG
```

| | |
|--------------------|--|
| Restriction Sites: | Sgfl-MluI |
| OTI Disclaimer: | Our molecular clone sequence data has been matched to the sequence identifier above as a point of reference. Note that the complete sequence of this clone is largely the same as the reference sequence but may contain minor differences , e.g., single nucleotide polymorphisms (SNPs). |
| Components: | The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The package also includes 100 pmols of both the corresponding 5' and 3' vector primers in separate vials. |



[View online »](#)

RefSeq: [NM_001504.2](#)

Summary: This gene encodes a G protein-coupled receptor with selectivity for three chemokines, termed CXCL9/Mig (monokine induced by interferon-g), CXCL10/IP10 (interferon-g-inducible 10 kDa protein) and CXCL11/I-TAC (interferon-inducible T cell a-chemoattractant). Binding of chemokines to this protein induces cellular responses that are involved in leukocyte traffic, most notably integrin activation, cytoskeletal changes and chemotactic migration. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. One of the isoforms (CXCR3-B) shows high affinity binding to chemokine, CXCL4/PF4 (PMID:12782716). [provided by RefSeq, Jun 2011]

Locus ID: 2833

MW: 16.6