

## Product datasheet for **SC206887**

### **KIR2DL2 (NM\_014219) Human 3' UTR Clone**

#### Product data:

**Product Type:** 3' UTR Clones  
**Product Name:** KIR2DL2 (NM\_014219) Human 3' UTR Clone  
**Vector:** pMirTarget (PS100062)  
**Symbol:** KIR2DL2  
**Synonyms:** CD158b; CD158B1; NKAT-6; NKAT6; p58.2  
**ACCN:** NM\_014219  
**Insert Size:** 541 bp  
**Insert Sequence:** >SC206887 3'UTR clone of NM\_014219  
The sequence shown below is from the reference sequence of NM\_014219. The complete sequence of this clone may contain minor differences, such as SNPs.  
**Blue**=Stop Codon **Red**=Cloning site

```
GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAAGCCAAGAAGGGCGGAAAGATCGCCGTG
TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC
TCCAGATCCAAAGTTGTCTCCTGCCCATGAGCACCACAGTCAGGCCTTGAGGGCGTCTTCTAGGGAGAC
AACAGCCCTGTCTCAAACCGGGTTGCCAGCTCCCATGTACCAGCAGCTGGAATCTGAAGGCATGAGTC
TGCATCTTAGGGCATCGCTCTTCTCACACCACAAATCTGAATGTGCCTCTCACTTGCTTACAAATGTC
TAAGGTCCCCACTGCCTGCTGGAGAAAAACACTCCTTTGCTTAGCCACAGTTCTCCATTTCACTT
GACCCCTGCCACCTCTCCAACCTAACTGGCTTACTTCCTAGTCTACTTGAGGCTGCAATCACACTGAG
GAACTCACAATCCAAACATACAAGAGGCTCCCTCTTAACGCAGCACTTAGACACGTGTTGTTCCACCT
TCCCTCATGCTGTTCCACCTCCCCTCAGACTAGCTTTCAGTCTTCTGTCAGCAGTAAAATTATATATT
TTTTAAAATAACTTCAATGTAGTTTTCCATCCTCAAATAAATCATGTCTGCCCCCATG
ACGCGTAAGCGGCCGCGCATCTAGATTGAAGAAAATGACCGACCAAGCGACGCCCAACCTGCCATCA
CGAGATTCGATTCCACCGCCGCTTCTATGAAAGG
```

**Restriction Sites:** SgfI-MluI

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the sequence identifier above as a point of reference. Note that the complete sequence of this clone is largely the same as the reference sequence but may contain minor differences, e.g., single nucleotide polymorphisms (SNPs).

**Components:** The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The package also includes 100 pmols of both the corresponding 5' and 3' vector primers in separate vials.

**RefSeq:** [NM\\_014219.2](#)



[View online >](#)

**Summary:**

Killer cell immunoglobulin-like receptors (KIRs) are transmembrane glycoproteins expressed by natural killer cells and subsets of T cells. The KIR genes are polymorphic and highly homologous and they are found in a cluster on chromosome 19q13.4 within the 1 Mb leukocyte receptor complex (LRC). The gene content of the KIR gene cluster varies among haplotypes, although several "framework" genes are found in all haplotypes (KIR3DL3, KIR3DP1, KIR3DL4, KIR3DL2). The KIR proteins are classified by the number of extracellular immunoglobulin domains (2D or 3D) and by whether they have a long (L) or short (S) cytoplasmic domain. KIR proteins with the long cytoplasmic domain transduce inhibitory signals upon ligand binding via an immune tyrosine-based inhibitory motif (ITIM), while KIR proteins with the short cytoplasmic domain lack the ITIM motif and instead associate with the TYRO protein tyrosine kinase binding protein to transduce activating signals. The ligands for several KIR proteins are subsets of HLA class I molecules; thus, KIR proteins are thought to play an important role in regulation of the immune response. [provided by RefSeq, Jul 2008]

**Locus ID:**

3803

**MW:**

19.8