

Product datasheet for SC206998

Synaptotagmin 3 (SYT3) (NM_032298) Human 3' UTR Clone

Product data:

Product Type: 3' UTR Clones
Product Name: Synaptotagmin 3 (SYT3) (NM_032298) Human 3' UTR Clone
Symbol: Synaptotagmin 3
Synonyms: SytIII
Mammalian Cell Selection: Neomycin
Vector: pMirTarget (PS100062)
ACCN: NM_032298
Insert Size: 538 bp
Insert Sequence: >SC206998 3'UTR clone of NM_032298
 The sequence shown below is from the reference sequence of NM_032298. The complete sequence of this clone may contain minor differences, such as SNPs.
 Blue=Stop Codon Red=Cloning site

```
GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAAGCCAAGAAGGGCGGAAAGATCGCCGTG
TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC
GGACTATCAGAGAAAGAGAACTCCGAGTGAGGGGTCTGGCCTAGGCCCGGGATCGGACCAGGCTCCCTC
AGGACCCATCCTTTCTGCCCGACCGTGAATTCATCTCCTTGAAGCCATAACGTCGAGCTGCTGGT
GCGGGGCAGCCCTGGCCCTAGGCTTCTAACCCCTGGAAGCGAGAGGATGAGAGGAGGCCGCCCAGCTC
CTTCTTTCAGGGTGGGGTCAATCAGCCTCCACTGTGTCTGTCTTTTCTCCCTGGGGCTCCCCCTCGA
GGCGAGGGGCCATGCATGTCTGGGGACCCCTGCCCCCAAAACCCTCTGTCTGTCTGTCTCTTTGC
TGTTTGTCCAAGACTCAGTGTCCCGACCCCTTGTCTCGCCGTGAATGTCAATGGGCAATCCTCTCTGT
CCTTTCAGACACACACACACTGTGTCCACCCCTTCTGTTCGCCACCCCTGCGTCTGGCCGGTCCCC
CCACTGTCTGTCTATCAACGCCAGAATAAACACACTCTGTGGGTCTCACTCAA
ACGCGTAAGCGGCCGCGGCATCTAGATTCAAGAAAATGACCGACCAAGCGACGCCCAACCTGCCATCA
CGAGATTCGATTCCACCGCCGCTTCTATGAAAGG
```

Restriction Sites: Sgfl-MluI
OTI Disclaimer: Our molecular clone sequence data has been matched to the sequence identifier above as a point of reference. Note that the complete sequence of this clone is largely the same as the reference sequence but may contain minor differences, e.g., single nucleotide polymorphisms (SNPs).



[View online »](#)

Components:	The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The package also includes 100 pmols of both the corresponding 5' and 3' vector primers in separate vials.
RefSeq:	NM_032298.3
Summary:	Ca(2+) sensor involved in Ca(2+)-dependent exocytosis of secretory vesicles through Ca(2+) and phospholipid binding to the C2 domain. Ca(2+) induces binding of the C2-domains to phospholipid membranes and to assembled SNARE-complexes; both actions contribute to triggering exocytosis (By similarity). Plays a role in dendrite formation by melanocytes (PubMed:23999003).[UniProtKB/Swiss-Prot Function]
Locus ID:	84258
MW:	19.4