

## Product datasheet for **SC207082**

### CD79B (NM\_000626) Human 3' UTR Clone

#### Product data:

**Product Type:** 3' UTR Clones  
**Product Name:** CD79B (NM\_000626) Human 3' UTR Clone  
**Vector:** pMirTarget (PS100062)  
**Symbol:** CD79B  
**Synonyms:** AGM6; B29; IGB  
**ACCN:** NM\_000626  
**Insert Size:** 516 bp  
**Insert Sequence:** >SC207082 3'UTR clone of NM\_000626  
The sequence shown below is from the reference sequence of NM\_000626. The complete sequence of this clone may contain minor differences, such as SNPs.  
**Blue**=Stop Codon **Red**=Cloning site

```
GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAAGCCAAGAAGGGCGGAAAGATCGCCGTG
TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC
TCTGTAGGTGAGCACCCAGGCCAGGAGTGAAGCCAGGTCGCCCCATGACCTGGGTGCAGGCTCCCTGG
CCTCAGTACTGCTTCGGAGCTGCCTGGCTCATGGCCCAACCCCTTTCTGGACCCCCAGCTGGCCTC
TGAAGCTGGCCCAACAGAGCTGCCATTTGTCTCCAGCCCTGGTCCCCAGCTCTTGCCAAAGGGCCTGG
AGTAGAAGGACAACAGGGCAGCAACTTGGAGGGAGTTCTCTGGGATGGACGGGACCCAGCCTTCTGGG
GGTGTATGAGGTGATCCGTCCCCACACATGGGATGGGGAGGCAGAGACTGGTCCAGAGCCCGCAAT
GGACTCGGAGCCGAGGGCCTCCCAGCAGAGCTTGGGAAGGGCCATGGACCCAACTGGGCCCCAGAAGAG
CCACAGGAACATCATTCTCTCCGCAACCACTCCCACCCAGGGAGGCCCTGGCCTCCAGTGCCTTCC
CCCGTGAATAAACGGTGTGTCTGAGAAACCA
ACGCGTAAGCGCCGCGCATCTAGATTGAAGAAAATGACCGACCAAGCGACGCCCAACCTGCCATCA
CGAGATTCGATTCCACCGCCGCTTCTATGAAAGG
```

**Restriction Sites:** SgfI-MluI

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the sequence identifier above as a point of reference. Note that the complete sequence of this clone is largely the same as the reference sequence but may contain minor differences, e.g., single nucleotide polymorphisms (SNPs).

**Components:** The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The package also includes 100 pmols of both the corresponding 5' and 3' vector primers in separate vials.

**RefSeq:** [NM\\_000626.4](#)



[View online >](#)

**Summary:** The B lymphocyte antigen receptor is a multimeric complex that includes the antigen-specific component, surface immunoglobulin (Ig). Surface Ig non-covalently associates with two other proteins, Ig-alpha and Ig-beta, which are necessary for expression and function of the B-cell antigen receptor. This gene encodes the Ig-beta protein of the B-cell antigen component. Alternatively spliced transcript variants encoding different isoforms have been described. [provided by RefSeq, Jul 2008]

**Locus ID:** 974

**MW:** 18