

## Product datasheet for **SC207350**

### **MED8 (NM\_052877) Human 3' UTR Clone**

#### Product data:

**Product Type:** 3' UTR Clones  
**Product Name:** MED8 (NM\_052877) Human 3' UTR Clone  
**Vector:** pMirTarget (PS100062)  
**Symbol:** MED8  
**Synonyms:** ARC32  
**ACCN:** NM\_052877  
**Insert Size:** 563 bp  
**Insert Sequence:** >SC207350 3'UTR clone of NM\_052877  
The sequence shown below is from the reference sequence of NM\_052877. The complete sequence of this clone may contain minor differences, such as SNPs.  
**Blue**=Stop Codon **Red**=Cloning site

```
GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAAGCCAAGAAGGGCGGAAAGATCGCCGTG
TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC
AAAAAAAAATGCCACCTGCAGCTCTGGTGAAGCTTGGGCCTGCTCTCCTTTCCATCCTCTAAGGAGCCA
ACTTGGCTTTTACCTGTCAAATAGTCATAAAGTCCCCTATCCTTTACCCACCTTATACACACGAGGCT
TTCTCAGGAAGTGGCTCTGCCAGGCAGGACTATGTGGAAAGGGTTTTTCCTTAGCACACGAAAAAGCC
CCTTCCCCTGGATTATGTTTCTATTTTGGAGGAGAAGGGAATTGCACTTCACACTGCCATCAGGGT
TTAGTTGACCTCATAATGGTGCCCACTTCTCGACTTTGGCCAGGATTTCTTCAAAGAAAAAGACTTT
CCTTCATTTCCCTAAGCCTGTGGCCAAATGGTGGACCAGAATGATGGTGGGAGGGGCAACCCCAAGT
AGCTTTGCCTGCTTTTATAAAGTTGAACAAATTGAATTTAGACATTCAGGCTAACCTGCCTTTCTAGT
ACTCCTTTGTTGGCATGGCAGGGGTTGAGTCAGCAGAAGTGGACCAAAGGATTCCTCTGAATAAAGTT
ATTTAAATTGA
ACGCGTAAGCGGCCGCGCATCTAGATTGAAGAAAATGACCGACCAAGCGACGCCAACCTGCCATCA
CGAGATTTGATTCCACCGCCGCTTCTATGAAAGG
```

**Restriction Sites:** SgfI-MluI

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the sequence identifier above as a point of reference. Note that the complete sequence of this clone is largely the same as the reference sequence but may contain minor differences, e.g., single nucleotide polymorphisms (SNPs).

**Components:** The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 µg dried plasmid DNA. The package also includes 100 pmols of both the corresponding 5' and 3' vector primers in separate vials.



[View online »](#)

RefSeq: [NM\\_052877.5](#)

**Summary:** This gene encodes a protein component of the mediator complex, which aids in transcriptional activation through interaction with RNA polymerase II and gene-specific transcription factors. The encoded protein may also function in ubiquitin ligation and protein degradation. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Jan 2013]

**Locus ID:** 112950

**MW:** 20.5