

## Product datasheet for SC207350

### MED8 (NM\_052877) Human 3' UTR Clone

#### Product data:

Product Type:	3' UTR Clones
Product Name:	MED8 (NM_052877) Human 3' UTR Clone
Symbol:	MED8
Synonyms:	ARC32
Mammalian Cell Selection:	Neomycin
Vector:	pMirTarget (PS100062)
ACCN:	NM_052877
Insert Size:	563 bp
Insert Sequence:	<p>&gt;SC207350 3'UTR clone of NM_052877</p> <p>The sequence shown below is from the reference sequence of NM_052877. The complete sequence of this clone may contain minor differences, such as SNPs.</p> <p>Blue=Stop Codon Red=Cloning site</p>

```

GGCAAGTTGGACGCCGCAAGATCCGCGAGATTCTCATTAAGGCCAAGAAGGGCGGAAAGATCGCCGTG
TAACAATTGGCAGAGCTCAGAATTCAAACGATCGCC
AAAAAAAAATGCCACCTGCAGCTCTGGTAGCTTGGGCTGCTCTCCTTTCCATCCTCTAAGGAGCCA
ACTTGGCTTTTACCTGTCAAATAGTCATAAAGTCCCTATCCTTTACCCACCTTATACACAGAGGCT
TTCTCAGGAAGTGGCTCTGCCAGGCAGGACTATGTGGGAAAGGGTTTTCTTAGCACACGAAAAAGCC
CCTTCCCCTGGATTCATGTTTCTTATTTTGGAGGGAGAAGGGAATTGCACTTCACACTGCCATCAGGGT
TTAGTTGACCTCATAATGGTGCCCACTTTCTCGACTTTGGCCAGGATTTCTTCAAAGAAAACGACTTT
CCTTCATTTCCCTAAGCCTGTGGCCCAATGGTGGACCAGAATGATGGTGGGAGGGGGCAACCCCAAGT
AGCTTTGCCTGCTTTTATAAAGTTGAACAAATTGAATTTAGACATTCAGGCTAACCTGCCTTTCTTAGT
ACTCCTTTGTTGGCATGGCAGGGGTTGAGTCAGCAGAAGTGGACCAAAGGATTCCTCTGAATAAAGTT
ATTAAATTGA
ACGCGTAAGCGGCCGCGGCATCTAGATTCTGAAGAAATGACCGACCAAGCGACGCCCAACCTGCCATCA
CGAGATTTGATTCCACCGCCGCTTCTATGAAAGG
  
```

Restriction Sites: SgfI-MluI

OTI Disclaimer: Our molecular clone sequence data has been matched to the sequence identifier above as a point of reference. Note that the complete sequence of this clone is largely the same as the reference sequence but may contain minor differences, e.g., single nucleotide polymorphisms (SNPs).


[View online »](#)

Components:	The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The package also includes 100 pmols of both the corresponding 5' and 3' vector primers in separate vials.
RefSeq:	<u><a href="#">NM_052877.5</a></u>
Summary:	This gene encodes a protein component of the mediator complex, which aids in transcriptional activation through interaction with RNA polymerase II and gene-specific transcription factors. The encoded protein may also function in ubiquitin ligation and protein degradation. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Jan 2013]
Locus ID:	112950
MW:	20.5