

## **Product datasheet for SC207825**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Metabotropic Glutamate Receptor 8 (GRM8) (NM 001127323) Human 3' UTR Clone

**Product data:** 

**Product Type:** 3' UTR Clones

Product Name: Metabotropic Glutamate Receptor 8 (GRM8) (NM\_001127323) Human 3' UTR Clone

Symbol: Metabotropic Glutamate Receptor 8
Synonyms: GLUR8; GPRC1H; mGlu8; MGLUR8

**Mammalian Cell** 

Selection:

Neomycin

**Vector:** pMirTarget (PS100062)

**ACCN:** NM\_001127323

**Insert Size:** 619 bp

Insert Sequence: >SC207825 3'UTR clone of NM\_001127323

The sequence shown below is from the reference sequence of NM\_001127323. The complete

sequence of this clone may contain minor differences, such as SNPs.

Blue=Stop Codon Red=Cloning site

GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAGGCCAAGAAGGGCGGAAAGATCGCCGTG

 ${\sf TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC}$ 

CGAGATTTCGATTCCACCGCCGCCTTCTATGAAAGG

Restriction Sites: Sgfl-Mlul

OTI Disclaimer: Our molecular clone sequence data has been matched to the sequence identifier above as a

point of reference. Note that the complete sequence of this clone is largely the same as the

reference sequence but may contain minor differences, e.g., single nucleotide

polymorphisms (SNPs).





## Metabotropic Glutamate Receptor 8 (GRM8) (NM\_001127323) Human 3' UTR Clone - SC207825

**Components:** The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The

package also includes 100 pmols of both the corresponding 5' and 3' vector primers in

separate vials.

**RefSeq:** NM 001127323.1

**Summary:** L-glutamate is the major excitatory neurotransmitter in the central nervous system and

activates both ionotropic and metabotropic glutamate receptors. Glutamatergic

neurotransmission is involved in most aspects of normal brain function and can be perturbed in many neuropathologic conditions. The metabotropic glutamate receptors are a family of G protein-coupled receptors, that have been divided into 3 groups on the basis of sequence homology, putative signal transduction mechanisms, and pharmacologic properties. Group I includes GRM1 and GRM5 and these receptors have been shown to activate phospholipase C. Group II includes GRM2 and GRM3 while Group III includes GRM4, GRM6, GRM7 and GRM8. Group II and III receptors are linked to the inhibition of the cyclic AMP cascade but differ in their agonist selectivities. Alternatively spliced transcript variants encoding different isoforms

have been described for this gene. [provided by RefSeq, Jul 2008]

Locus ID: 2918 MW: 23.6