

Product datasheet for SC208106

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

p19 INK4d (CDKN2D) (NM_079421) Human 3' UTR Clone

Product data:

Product Type: 3' UTR Clones

Product Name: p19 INK4d (CDKN2D) (NM 079421) Human 3' UTR Clone

Symbol: p19 INK4d

Synonyms: INK4D; p19; p19-INK4D

Mammalian Cell

Selection:

Neomycin

..........

Vector:

pMirTarget (PS100062)

ACCN: NM 079421

Insert Size:

628 bp

Insert Sequence:

>SC208106 3'UTR clone of NM_079421

The sequence shown below is from the reference sequence of NM_079421. The complete

sequence of this clone may contain minor differences, such as SNPs.

Blue=Stop Codon Red=Cloning site

GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAGGCCAAGAAGGGCGGAAAGATCGCCGTG

TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC

GATCACA

CGAGATTTCGATTCCACCGCCGCCTTCTATGAAAGG

Restriction Sites: Sgfl-Mlul

OTI Disclaimer: Our molecular clone sequence data has been matched to the sequence identifier above as a

point of reference. Note that the complete sequence of this clone is largely the same as the

reference sequence but may contain minor differences, e.g., single nucleotide

polymorphisms (SNPs).





p19 INK4d (CDKN2D) (NM_079421) Human 3' UTR Clone - SC208106

Components: The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The

package also includes 100 pmols of both the corresponding 5' and 3' vector primers in

separate vials.

RefSeq: <u>NM 079421.3</u>

Summary: The protein encoded by this gene is a member of the INK4 family of cyclin-dependent kinase

inhibitors. This protein has been shown to form a stable complex with CDK4 or CDK6, and prevent the activation of the CDK kinases, thus function as a cell growth regulator that controls cell cycle G1 progression. The abundance of the transcript of this gene was found to oscillate in a cell-cycle dependent manner with the lowest expression at mid G1 and a maximal expression during S phase. The negative regulation of the cell cycle involved in this

protein was shown to participate in repressing neuronal proliferation, as well as

spermatogenesis. Two alternatively spliced variants of this gene, which encode an identical

protein, have been reported. [provided by RefSeq, Jul 2008]

Locus ID: 1032 **MW:** 23.9