

## **Product datasheet for SC208235**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Sumo 1 (SUMO1) (NM\_001005781) Human 3' UTR Clone

**Product data:** 

**Product Type:** 3' UTR Clones

Product Name: Sumo 1 (SUMO1) (NM\_001005781) Human 3' UTR Clone

**Vector:** pMirTarget (PS100062)

Symbol: SUMO1

Synonyms: DAP1; GMP1; OFC10; PIC1; SENP2; SMT3; SMT3C; SMT3H3; UBL1

ACCN: NM\_001005781

**Insert Size:** 640 bp

Insert Sequence: >SC208235 3' UTR clone of NM\_001005781

The sequence shown below is from the reference sequence of NM\_001005781. The complete sequence of this clone may contain minor differences, such as SNPs. Red=Cloning site

Blue=Stop Codon

CAATTGGCAGAGCTCAGAATTCAAGCGATCGC

GAATAAATGG

**ACGCGT**AAGCGGCCGCGGCATCTAGATTCGAAGAAAATGACCG

**Restriction Sites:** Sgfl-Mlul

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the sequence identifier above as a

point of reference. Note that the complete sequence of this clone is largely the same as the

reference sequence but may contain minor differences, e.g., single nucleotide

polymorphisms (SNPs).





## Sumo 1 (SUMO1) (NM\_001005781) Human 3' UTR Clone - SC208235

**Components:** The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The

package also includes 100 pmols of both the corresponding 5' and 3' vector primers in

separate vials.

RefSeq: <u>NM 001005781.1</u>

**Summary:** This gene encodes a protein that is a member of the SUMO (small ubiquitin-like modifier)

protein family. It functions in a manner similar to ubiquitin in that it is bound to target proteins as part of a post-translational modification system. However, unlike ubiquitin which targets proteins for degradation, this protein is involved in a variety of cellular processes, such as nuclear transport, transcriptional regulation, apoptosis, and protein stability. It is not active until the last four amino acids of the carboxy-terminus have been cleaved off. Several pseudogenes have been reported for this gene. Alternate transcriptional splice variants encoding different isoforms have been characterized. [provided by RefSeq, Jul 2008]

**Locus ID:** 7341