

Product datasheet for SC208645

E2F5 (NM_001083588) Human 3' UTR Clone

Product data:

Product Type: 3' UTR Clones

Product Name: E2F5 (NM_001083588) Human 3' UTR Clone

Symbol: E2F5
Synonyms: E2F-5

Mammalian Cell Neomycin

Selection:

Vector: pMirTarget (PS100062)

ACCN: NM_001083588

Insert Size: 686 bp

Insert Sequence: >SC208645 3'UTR clone of NM_001083588

The sequence shown below is from the reference sequence of NM_001083588. The complete

sequence of this clone may contain minor differences, such as SNPs.

Blue=Stop Codon Red=Cloning site

GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAGGCCAAGAAGGGCGGAAAGATCGCCGTG

TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC

CGAGATTTCGATTCCACCGCCGCCTTCTATGAAAGG

Restriction Sites: Sgfl-Mlul

OTI Disclaimer: Our molecular clone sequence data has been matched to the sequence identifier above as a

point of reference. Note that the complete sequence of this clone is largely the same as the

reference sequence but may contain minor differences, e.g., single nucleotide

polymorphisms (SNPs).



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



E2F5 (NM_001083588) Human 3' UTR Clone - SC208645

Components: The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The

package also includes 100 pmols of both the corresponding 5' and 3' vector primers in

separate vials.

RefSeq: <u>NM 001083588.2</u>

Summary: The protein encoded by this gene is a member of the E2F family of transcription factors. The

E2F family plays a crucial role in the control of cell cycle and action of tumor suppressor proteins and is also a target of the transforming proteins of small DNA tumor viruses. The E2F proteins contain several evolutionarily conserved domains that are present in most members of the family. These domains include a DNA binding domain, a dimerization domain which determines interaction with the differentiation regulated transcription factor proteins (DP), a transactivation domain enriched in acidic amino acids, and a tumor suppressor protein association domain which is embedded within the transactivation domain. This protein is differentially phosphorylated and is expressed in a wide variety of human tissues. It has higher identity to E2F4 than to other family members. Both this protein

and E2F4 interact with tumor suppressor proteins p130 and p107, but not with pRB.

Alternative splicing results in multiple variants encoding different isoforms. [provided by

RefSeq, Jul 2008]

Locus ID: 1875

MW: 26.4