

Product datasheet for SC208795

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Mortality Factor 4 like 2 (MORF4L2) (NM_001142424) Human 3' UTR Clone

Product data:

Product Type: 3' UTR Clones

Product Name: Mortality Factor 4 like 2 (MORF4L2) (NM_001142424) Human 3' UTR Clone

Vector: pMirTarget (PS100062)

Symbol: MORF4L2

Synonyms: MORFL2; MRGX ACCN: NM_001142424

Insert Size: 693 bp

Insert Sequence: >SC208795 3'UTR clone of NM_001142424

The sequence shown below is from the reference sequence of NM_001142424. The complete

sequence of this clone may contain minor differences, such as SNPs.

Blue=Stop Codon Red=Cloning site

GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAGGCCAAGAAGGGCGGAAAGATCGCCGTG

TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC

CGAGATTTCGATTCCACCGCCGCCTTCTATGAAAGG

Restriction Sites: Sgfl-Mlul

OTI Disclaimer: Our molecular clone sequence data has been matched to the sequence identifier above as a

point of reference. Note that the complete sequence of this clone is largely the same as the

reference sequence but may contain minor differences, e.g., single nucleotide

polymorphisms (SNPs).





Mortality Factor 4 like 2 (MORF4L2) (NM_001142424) Human 3' UTR Clone - SC208795

Components: The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The

package also includes 100 pmols of both the corresponding 5' and 3' vector primers in

separate vials.

RefSeq: <u>NM 001142424.2</u>

Summary: Component of the NuA4 histone acetyltransferase complex which is involved in

transcriptional activation of select genes principally by acetylation of nucleosomal histone H4

and H2A. This modification may both alter nucleosome - DNA interactions and promote

interaction of the modified histones with other proteins which positively regulate

transcription. This complex may be required for the activation of transcriptional programs associated with oncogene and proto-oncogene mediated growth induction, tumor suppressor mediated growth arrest and replicative senescence, apoptosis, and DNA repair. The NuA4 complex ATPase and helicase activities seem to be, at least in part, contributed by the association of RUVBL1 and RUVBL2 with EP400. NuA4 may also play a direct role in DNA repair when directly recruited to sites of DNA damage. Also component of the MSIN3A complex which acts to repress transcription by deacetylation of nucleosomal histones.

[UniProtKB/Swiss-Prot Function]

Locus ID: 9643 **MW:** 27.1