

## **Product datasheet for SC208819**

LILRB1 (NM 001081639) Human 3' UTR Clone

## **Product data:**

**Product Type:** 3' UTR Clones

Product Name: LILRB1 (NM\_001081639) Human 3' UTR Clone

Symbol: LILRB1

Synonyms: CD85J; ILT-2; ILT2; LIR-1; LIR1; MIR-7; MIR7; PIR-B; PIRB

**Mammalian Cell** 

Selection:

Neomycin

**Vector:** pMirTarget (PS100062)

**ACCN:** NM\_001081639

**Insert Size:** 708 bp

Insert Sequence: >SC208819 3'UTR clone of NM\_001081639

The sequence shown below is from the reference sequence of NM\_001081639. The complete

sequence of this clone may contain minor differences, such as SNPs.

Blue=Stop Codon Red=Cloning site

GGCAAGTTGGACGCCCGCAAGATCCGCGAGATTCTCATTAAGGCCAAGAAGGGCGGAAAGATCGCCGTG

 ${\sf TAACAATTGGCAGAGCTCAGAATTCAAGCGATCGCC}$ 

AAAAAAAGAAAGAAAAA

**ACGCGT**AAGCGGCCGCGCATCTAGATTCGAAGAAAATGACCGACCAAGCGACGCCCAACCTGCCATCA

CGAGATTTCGATTCCACCGCCGCCTTCTATGAAAGG

**Restriction Sites:** Sgfl-Mlul

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the sequence identifier above as a

point of reference. Note that the complete sequence of this clone is largely the same as the

reference sequence but may contain minor differences, e.g., single nucleotide

polymorphisms (SNPs).



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## LILRB1 (NM\_001081639) Human 3' UTR Clone - SC208819

**Components:** The cDNA clone is shipped in a 2-D bar-coded Matrix tube as 10 ug dried plasmid DNA. The

package also includes 100 pmols of both the corresponding 5' and 3' vector primers in

separate vials.

**RefSeq:** <u>NM 001081639.4</u>

**Summary:** This gene is a member of the leukocyte immunoglobulin-like receptor (LIR) family, which is

found in a gene cluster at chromosomal region 19q13.4. The encoded protein belongs to the subfamily B class of LIR receptors which contain two or four extracellular immunoglobulin domains, a transmembrane domain, and two to four cytoplasmic immunoreceptor tyrosine-based inhibitory motifs (ITIMs). The receptor is expressed on immune cells where it binds to MHC class I molecules on antigen-presenting cells and transduces a negative signal that inhibits stimulation of an immune response. It is thought to control inflammatory responses and cytotoxicity to help focus the immune response and limit autoreactivity. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by

RefSeq, Jul 2008]

**Locus ID:** 10859

MW: 26.2